

Clinical pharmacists can aid patients with uncontrolled T2DM

28 October 2013



Clinical pharmacists can play a role in helping patients with poorly controlled type 2 diabetes, according to a case study published in the October issue of *Clinical Diabetes*.

(HealthDay)—Clinical pharmacists can play a role in helping patients with poorly controlled type 2 diabetes, according to a case study published in the October issue of *Clinical Diabetes*.

Travis E. Sonnett, Pharm.D., from Washington State University, and Tyler Galloway, Pharm.D., from the Mann-Grandstaff Veteran Affairs Medical Center—both in Spokane, examined the impact of a clinical pharmacy specialist on management of a wheelchair-bound, 64-year-old man with a history of uncontrolled [type 2 diabetes](#) (hemoglobin A1c [A1c], 10.0 percent). The patient reported poor adherence to his insulin regimen, but close adherence to his oral medication regimen. He experienced progression of peripheral neuropathy and retinopathy.

The authors reported that the clinical pharmacist used a motivational interview technique to discuss the patient's failure to follow the current insulin regimen and poor eating. They developed a plan to monitor progress and agreed on an initial goal of A1c

APA citation: Clinical pharmacists can aid patients with uncontrolled T2DM (2013, October 28) retrieved 29 September 2022 from <https://medicalxpress.com/news/2013-10-clinical-pharmacists-aid-patients->

[uncontrolled.html](#)

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.