

Effect of reducing blood pressure with medications immediately following ischemic stroke

17 November 2013

Jiang He, M.D., Ph.D., of the Tulane University School of Public Health and Tropical Medicine, New Orleans, and colleagues examined whether moderate lowering of blood pressure within the first 48 hours after the onset of an acute ischemic stroke would reduce death and major disability at 14 days or hospital discharge.

"Stroke is the second leading cause of death and the leading cause of serious, long-term disability worldwide. Clinical trials have documented that lowering [blood pressure](#) reduces the risk of [stroke](#) in hypertensive and normotensive patients with a history of stroke or [transient ischemic attack](#). Although the benefit of lowering blood pressure for primary and secondary prevention of stroke has been established, the effect of immediate antihypertensive treatment in patients with acute ischemic stroke and elevated blood pressure is uncertain," according to background information in the article.

The China Antihypertensive Trial in Acute Ischemic Stroke, a randomized controlled trial, was conducted among 4,071 patients with ischemic stroke within 48 hours of symptom onset and elevated [systolic blood pressure](#). Patients were recruited from 26 hospitals across China between August 2009 and May 2013. Patients (n = 2,038) were assigned to receive antihypertensive treatment (aimed at lowering systolic blood pressure by 10 percent to 25 percent within the first 24 hours after randomization, achieving blood pressure less than 140/90 mm Hg within 7 days, and maintaining this level during hospitalization) or to discontinue all antihypertensive medications (control) during hospitalization (n = 2,033).

Average systolic blood pressure was reduced from 166.7 mm Hg to 144.7 mm Hg (-12.7 percent) within 24 hours in the antihypertensive treatment

group and from 165.6 mm Hg to 152.9 mm Hg (-7.2 percent) in the control group within 24 hours after randomization. Average systolic blood pressure was 137.3 mm Hg in the antihypertensive treatment group and 146.5 mm Hg in the control group at day 7 after randomization. The primary outcome (a combination of death and major disability at 14 days or hospital discharge) did not differ between treatment groups (683 events [antihypertensive treatment] vs. 681 events [control]) at 14 days or [hospital discharge](#). The secondary composite outcome of death and major disability at 3-month posttreatment follow-up did not differ between treatment groups.

These findings suggest that the decision to lower blood pressure with antihypertensive treatment in patients with [acute ischemic stroke](#) does not improve or worsen outcome and therefore should be based on individual clinical judgment, the authors write.

More information: *JAMA*
doi:10.1001/jama.2013.282543

Provided by The JAMA Network Journals

APA citation: Effect of reducing blood pressure with medications immediately following ischemic stroke (2013, November 17) retrieved 2 June 2022 from <https://medicalxpress.com/news/2013-11-effect-blood-pressure-medications-immediately.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.