

# Pre-op MRI of margins predicts rectal cancer outcomes

8 December 2013



Committee on Cancer TNM-based criteria for assessing risk of LR, DFS, and OS," the authors write.

**More information:** [Abstract](#)  
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2013 [HealthDay](#). All rights reserved.

(HealthDay)—Preoperative high-resolution magnetic resonance imaging (MRI) assessment of circumferential resection margin (CRM) status predicts survival and other disease progression outcomes in patients with rectal cancer, according to a study published online Nov. 25 in the *Journal of Clinical Oncology*.

Fiona G.M. Taylor, from Mayday University Hospital in Croydon, U.K., and colleagues conducted a follow-up study in 374 patients with [rectal cancer](#). Preoperative high-resolution pelvic MRIs were used to assess CRM, and the relationship of MRI assessment and outcomes were evaluated.

The researchers found that five-year overall survival (OS) was 62.2 percent in patients with MRI-clear CRM compared to 42.2 percent in patients with MRI-involved CRM (hazard ratio, 1.97). Disease-free survival (DFS) at five years was 67.2 percent for MRI-clear CRM versus 47.3 percent for MRI-involved CRM (hazard ratio, 1.65). For MRI-involved CRM, local recurrence (LR) was 3.50. In multivariate analysis, MRI-involved CRM was the only preoperative staging parameter that remained significant for OS, DFS, and LR.

"High-resolution MRI preoperative assessment of CRM status is superior to American Joint

APA citation: Pre-op MRI of margins predicts rectal cancer outcomes (2013, December 8) retrieved 19 June 2022 from <https://medicalxpress.com/news/2013-12-pre-op-mri-margins-rectal-cancer.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*