

Infectious diarrhea germs stick to healthcare worker hands

December 23 2013

A new study finds nearly one in four healthcare workers' hands were contaminated with *Clostridium difficile* spores after routine care of patients infected with the bacteria. The study was published in the January issue of *Infection Control and Hospital Epidemiology*, the journal of the Society for Healthcare Epidemiology of America.

"This is the first known study focusing on the carriage of viable *C. difficile* spores on [healthcare workers](#) hands," said Caroline Landelle, PharmD, PhD, lead author of the study. "Because *C. difficile* spores are so resistant and persistent to disinfection, glove use is not an absolute barrier against the contamination of healthcare workers' hands. Effective [hand](#) hygiene should be performed, even in non-outbreak settings."

Researchers compared hand contamination rates among healthcare workers caring for patients with *C. difficile* with healthcare workers caring for non-colonized patients after routine patient care and before hand hygiene. All patients with *C. difficile* were being treated with [infection control](#) measures that consisted of (1) placing patients into a single-bed room with dedicated equipment; (2) wearing disposable gowns with full-length sleeves and a pair of gloves on entering the room; (3) [hand hygiene](#) with alcohol-based hand rub before wearing gloves, before and after body fluid exposure, and hand washing with medicated soap and water followed by use of alcohol-based hand rub after glove removal; and (4) daily room cleaning with a hypochlorite-based disinfectant.

Contamination of healthcare workers' hands occurred with high-risk contact (e.g., patient washing, digital rectal exam, bed linen change, colonoscopy) or when workers didn't use gloves. Hand contamination was also associated with the duration of high-risk contact and was more common among nursing assistants (42 percent) than among other healthcare workers (19 percent for nurses and 23 percent for physicians), likely because nursing assistants had more high-risk contact.

In a commentary accompanying the study, author Aurora Pop-Vicas, MD, notes that "this study offers a vivid insight into why *C. difficile* might be so stubbornly persistent in our hospital...much work remains to be done in implementing what is known about the prevention of the spread of this bacteria through horizontal transmission. Additional measures include improvement in antimicrobial stewardship programs and effective environmental cleaning within healthcare institutions."

The number of patients hospitalized with a primary *C. difficile* diagnosis in the United States more than tripled during 2000-2009. Key risk factors of contracting *C. difficile* include prior receipt of antibiotic therapy, age, severity of underlying disease, length of hospital stay and prior occupancy of patient rooms. However, many healthcare workers may be passing on this highly contagious bacteria to patients even after routine alcohol-based hand rubbing. This points to the need for routine hand washing with soap and water, rather than alcohol-based hand rub, after care of *C. difficile* patients in all settings.

In the Compendium of Strategies to Prevent Healthcare-Associated Infections, SHEA recommends that health professionals clean hands with soap and water after caring for [patients](#) with *C. difficile* infections. Evidence shows that soap and water is superior to alcohol-based sanitizers for removing *C. difficile* spores. SHEA plans to release an updated Compendium in 2014.

More information: C. Landelle, M. Verachten, P. Legrand E. Girou, F. Barbut, C. Brun Buisson. "Contamination of Healthcare Workers' Hands with Clostridium difficile Spores after Caring for Patients with C. difficile Infection." *Infection Control and Hospital Epidemiology* 35:1 (January 2014).

Provided by Society for Healthcare Epidemiology of America

Citation: Infectious diarrhea germs stick to healthcare worker hands (2013, December 23)
retrieved 21 November 2023 from

<https://medicalxpress.com/news/2013-12-infectious-diarrhea-germs-healthcare-worker.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.