

Carotid Artery MRI helps predict likelihood of strokes and heart attacks

4 March 2014



This shows a transverse (a) T2-weighted, (b) unenhanced T1-weighted, and (c) gadolinium-enhanced T1-weighted MR images in an 82-year-old man. Pulse sequence parameters are described in Table 1. The left ICA wall shows no plaque or significant wall thickening. The right ICA shows plaque with a lipid core (arrow). ECA = external carotid artery, IJ = internal jugular vein. Credit: Radiological Society of North America


Noninvasive imaging of carotid artery plaque with MRI can accurately predict future cardiovascular events like strokes and heart attacks in people without a history of cardiovascular disease, according to a new study published online in the journal *Radiology*.

Researchers have long known that some arterial plaque is more dangerous because of its vulnerability to rupture. MRI can discern features of vulnerable plaque, such as a lipid core with a thin fibrous cap. This ability makes MRI a potentially valuable tool for identifying patients at risk for subsequent cardiovascular events.

To study the predictive value of MRI plaque imaging, researchers performed [carotid artery](#) ultrasound and MRI on 946 asymptomatic patients from the Multi-Ethnic Study of Atherosclerosis (MESA). The carotid arteries are the large vessels located on each side of the neck that carry oxygenated blood to the front part of the brain. They are highly accessible for imaging, and their condition tends to mirror that of the coronary arteries that supply the heart with oxygenated blood.

The researchers used ultrasound to assess carotid wall thickness and MRI to define carotid plaque composition and the remodeling index, a measure of changes in vessel size. Imaging results were compared with cardiovascular events, including heart attacks, stroke and death, for an average of 5.5 years after examination.

"We studied asymptomatic individuals with a low risk of cardiovascular events at baseline and used noninvasive imaging to predict the risk of an event downstream," said David A. Bluemke, M.D, Ph.D., from the National Institutes of Health Clinical Center in Bethesda, Md. "This is the first population-based prospective study to determine if vulnerable plaque features by MRI add to the risk of a cardiovascular event beyond the traditional risk factors."



This shows transverse gadolinium-enhanced T1-weighted MR images obtained superior to the carotid artery bifurcation in a 72-year-old man. L = ICA lumen. (a) Low-signal-intensity calcium (arrow) and lipid core (arrowheads) can be seen. (b) Note contouring of the ICA. The outer adventitial wall (red), lipid core (blue), calcification (green), and vessel lumen (purple) are visible. Credit: Radiological Society of North America

Cardiovascular events occurred in 59 of the patients. Abnormal thickening of the carotid artery

wall and the presence of a lipid core and calcium in the internal carotid artery on MRI were significant predictors of subsequent events. A lipid core was present in almost half of the patients who had an event, compared with only 17.8 percent of those who did not have an event.

"The primary factors that predicted future risk were measures of vessel wall thickness in combination with the presence or absence of a lipid core," Dr. Bluemke said. "The presence of a lipid core was 50 percent more common in people who had subsequent events."

Use of MRI improved the reclassification of baseline cardiovascular risk in the study group. When both the carotid remodeling index and lipid core were used for risk stratification, approximately 16 percent more patients with events and 7 percent without events were correctly reclassified compared with the use of traditional risk factors.

"The results bolster the use of MRI as a surrogate marker of efficacy in therapeutic studies and point to a role in determining which patients might need more aggressive treatments," Dr. Bluemke said. "As risk factor prediction gets better, we'll be able to screen more intelligently and use more intensive treatments in those individuals who face a higher risk of [cardiovascular events](#)."


This diagram shows representative changes in carotid wall area, lumen area, and remodeling index for the ICA and CCA. Representative values of lumen and wall areas are taken from Table 1. The normal remodeling index tends to be approximately 0.4 independent of vessel size. The remodeling index increases with relatively increased atherosclerotic burden. Diagrams are not to scale. Credit: Radiological Society of North America

Dr. Bluemke noted that sequences for plaque composition could be readily added to existing carotid MRI angiography protocols for clinical purposes.

"Carotid MRI has significant advantages," he said. "The results are more reproducible than those of ultrasound."

Availability and cost effectiveness could limit the use of MRI, Dr. Bluemke cautioned. However, ultrasound and computed tomography (CT) are viable alternatives for screening, he said, especially with improvements in CT dose reduction.

More information: "Carotid Artery Plaque Morphology and Composition in Relation to Incident Cardiovascular Events: The Multi-Ethnic Study of Atherosclerosis (MESA)." *Radiology*, 2014.



Provided by Radiological Society of North America

APA citation: Carotid Artery MRI helps predict likelihood of strokes and heart attacks (2014, March 4) retrieved 3 May 2021 from <https://medicalxpress.com/news/2014-03-carotid-artery-mri-likelihood-heart.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.