

# Nocturnal respiratory rate predicts cardiac risk after MI

March 10 2014

---



(HealthDay)—Among survivors of acute myocardial infarction (MI), nocturnal respiratory rate (NRR) is significantly associated with cardiac mortality, particularly non-sudden cardiac death, according to research published online March 5 in the *Journal of the American College of Cardiology*.

Michael Dommasch, M.D., of the Medizinische Klinik der Technischen Universität München, and colleagues applied an algorithm to data from Holter electrocardiographic recordings for 1,538 MI survivors to determine NRR. The researchers examined the association between NRR and cardiac death.

The researchers found that NRR was 18.6 per minute or greater in 384 patients and less than 18.6 per minute in 1,154 patients; five-year [cardiac mortality](#) in these groups was 13.7 and 3.3 percent, respectively (P

sudden cardiac death (hazard ratio, 4.56; 95 percent confidence interval, 2.21 to 9.43; P

"Among patients with left ventricular ejection fraction  $\leq 35$  percent, i.e., [implantable cardioverter defibrillator](#) (ICD) candidates, increased NRR identifies a group of patients with frequent non-sudden cardiac death, i.e., a mode of death likely not ICD preventable," the authors write.

"Independent validation of this new risk stratification approach is necessary before it can be incorporated into clinical decision-making."

**More information:** [Full Text \(subscription or payment may be required\)](#)

Copyright © 2014 [HealthDay](#). All rights reserved.

Citation: Nocturnal respiratory rate predicts cardiac risk after MI (2014, March 10) retrieved 1 February 2023 from <https://medicalxpress.com/news/2014-03-nocturnal-respiratory-cardiac-mi.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.