


Neck circumference not tied to subclinical atherosclerosis

27 March 2014

[Full Text \(subscription or payment may be required\)](#)



Copyright © 2014 [HealthDay](#). All rights reserved.

(HealthDay)—Neck circumference (NC) is not associated with either coronary or carotid subclinical atherosclerosis, according to a study published online March 19 in *Clinical Cardiology*.

Yashashwi Pokharel, M.D., from the Baylor College of Medicine in Houston, and colleagues measured NC (midway between the midcervical spine and midanterior neck) in 845 retired National Football League (NFL) players. Carotid ultrasound was used to identify [carotid artery](#) plaque (CAP).

The researchers found that 21 percent of participants had metabolic syndrome (MetS), while 62 and 56 percent, respectively, had [coronary artery calcium](#) (CAC) and CAP. Participants with MetS had a significantly higher median NC compared to those without MetS (P body mass index, results were similar).

"In retired NFL players with a high prevalence of CAC and CAP, NC was not associated with coronary or carotid subclinical atherosclerosis," the authors write. "NC may not be the most appropriate risk marker for atherosclerosis."

Several authors disclosed financial ties to the pharmaceutical and medical device industries.

More information: [Abstract](#)

APA citation: Neck circumference not tied to subclinical atherosclerosis (2014, March 27) retrieved 3 October 2022 from <https://medicalxpress.com/news/2014-03-neck-circumference-tied-subclinical-atherosclerosis.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.