

Two new genes linked to intellectual disability

31 March 2014

Researchers at the Centre for Addiction and Mental Health have discovered two new genes linked to intellectual disability, according to two research studies published concurrently this month in the journals *Human Genetics* and *Human Molecular Genetics*.

"Both studies give clues to the different pathways involved in normal neurodevelopment," says CAMH Senior Scientist Dr. John Vincent, who heads the MiND (Molecular Neuropsychiatry and Development) Laboratory in the Campbell Family Mental Health Research Institute at CAMH. "We are building up a body of knowledge that is informing us which kinds of [genes](#) are important to, and involved in, [intellectual disabilities](#)."

In the first study, Dr. Vincent and his team used microarray genotyping to map the genes of a large Pakistani family which had intermarriage. Five members of the youngest generation were affected with mild to moderate intellectual disability. Dr. Vincent identified a truncation in the *FBXO31* gene, which plays a role in the way that proteins are processed during development of neurons, particularly in the cerebellar cortex.

In the second study, using the same techniques, Dr. Vincent and his team analyzed the genes of two families with intermarriage, one Austrian and one Pakistani, and identified a disruption in the *METTL23* gene linked to mild recessive intellectual disability. The *METTL23* gene is involved in methylation—a process important to brain development and function.

About one per cent of children worldwide are affected by non-syndromic (i.e., the absence of any other clinical features) intellectual disability, a condition characterized by an impaired capacity to learn and process new or complex information, leading to decreased cognitive functioning and social adjustment. Although trauma, infection and external damage to the unborn fetus can lead to an

intellectual disability, genetic defects are a principal cause.

These studies were part of an ongoing study of affected families in Pakistan, where the cultural tradition of large families and consanguineous (inter-) marriages among first cousins increases the likelihood of inherited intellectual disability in offspring.

"Although it is easier to find and track genes in consanguineous families, these genes are certainly not limited to them," Dr. Vincent points out. A recent study estimated that 13

Provided by Centre for Addiction and Mental Health

APA citation: Two new genes linked to intellectual disability (2014, March 31) retrieved 28 June 2022 from <https://medicalxpress.com/news/2014-03-genes-linked-intellectual-disability.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.