

HIV-positive children more likely to develop drug resistance

28 May 2014, by Keith Brannon



The study emphasizes the importance of drug adherence for children and adolescents with HIV, says Dr. Russell Van Dyke. Credit: Paula Burch-Celentano

Children born with HIV are far more likely to develop resistance to antiretroviral drugs than adults, according to a new Tulane University School of Medicine study.

Researchers following almost 450 children enrolled in the Pediatric HIV/AIDS Cohort Study, one of the largest studies of HIV-positive children in the United States, found that 74 percent had developed resistance to at least one form of drug treatment and 30 percent were resistant to at least two classes of HIV treatment drugs. In comparison, nearly 36 percent of adults with HIV have resistance to one form of [drug treatment](#) and only 12 percent have resistance to at least two classes.

The study, which tracks patients at 14 sites across the country, followed participants from 7 to 16 years of age at enrollment.

"The problem with [drug resistance](#) is that once you develop it, it never goes away," said principal investigator Dr. Russell Van Dyke, professor of

pediatric infectious diseases. "Some patients with very resistant virus have no effective treatment options. Resistant virus is the major reason for death among youth with perinatal HIV."

Fortunately, most adolescents with resistant HIV remain sensitive to newer agents from all classes, allowing salvage therapy. Just one child in the study had resistance to all HIV drugs and only 18 percent had resistance to one drug from each of the three primary classes of HIV medications.

The study emphasizes the importance of drug adherence for children and adolescents with HIV who typically take multiple medications daily to manage the disease. Once daily, single-pill HIV medications for adults are not yet available for children, Van Dyke said.

"The best way to prevent resistance from developing is to take your medicine and suppress your viral load," Van Dyke said. "You develop resistance when you take some of your medications but not all. Then you've got virus that is replicating in the face of taking your medication. Lack of adherence is the major reason [resistance](#) develops."

The study abstract was presented recently at the Conference on Retroviruses and Opportunistic Infections.

More information: The abstract is available online: croiconference.org/sites/all/abstracts/897.pdf

Provided by Tulane University

APA citation: HIV-positive children more likely to develop drug resistance (2014, May 28) retrieved 28 April 2021 from <https://medicalxpress.com/news/2014-05-hiv-positive-children-drug-resistance.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.