

# New research shows parental presence improves the quality of child anaesthesia

June 1 2014

---

New research presented at this year's Euroanaesthesia meeting in Stockholm shows that having parents present during the induction of their child's anaesthesia improves the quality of that anaesthesia. The study is by Dr Alicia Sánchez, Department of Anaesthesiology, Hospital de Sagunto, Spain, and colleagues.

Induction of [anaesthesia](#) can be distressing for [parents](#) and [children](#). The effect of parental presence at anaesthesia induction on children anxiety and children anaesthesia compliance has been previously investigated but the few studies to date have produced contradictory results; and nobody has investigated issues around parental experience and total perceived quality. The aim of this randomised controlled trial was to evaluate the impact of parental presence during induction of anaesthesia on parent and children anxiety during the perioperative period and perceived quality.

Nowadays, common practice in most hospitals consists of children being separated from their parents at the entrance on the [operating room](#)," says Dr Sánchez. "However some hospitals allow one of the parents to stay with their child after surgery, in the post-anaesthetic care unit."

Local ethics committee approval was granted, and the researchers then obtained written informed consent from the parents of 60 child patients scheduled for tonsillectomy. Those who had previously had surgery were excluded. Patients were randomly allocated to two groups: one with the presence of a children's parent during induction of anaesthesia (P) or

children who were separated from their parents at the entrance to the operating room (the usual practice) and thus the control group (C). 30 children were allocated to each group.

Analysis of parental anxiety was carried out using the State Trait Anxiety Inventory (STAI), a parental self-report measuring state of mind and trait anxiety. The form was filled in the waiting area (baseline and moment 1); after separation (C group) (moment 2) or just before entering the operating room (P group) (moment 2); during induction of anaesthesia (moment 3); and in the postanesthetic care unit (moment 4).

Another anaesthetist used the Modified Yale Preoperative Anxiety Scale (mYPAS) to measure children's anxiety at the same moments. mYPAS scores >40 were classified as anxious.

The Induction Compliance Checklist (ICC) represents the negative behaviours present during induction, with an ICC score of 4 or more considered poor behavioural compliance. This was completed by an anaesthetist; thus two anaesthetists were present during the study, one to conduct the anaesthesia and one to complete the paperwork and ensure the correct procedures were being followed for the trial. Finally, when leaving the post-anaesthetic care unit all parents answered a questionnaire about their experience and evaluated the experience in a scale from 0 to 10.

All children and parents completed the study. The researchers found that [anxiety](#) scores were much lower in group P than in group C, with the scores in group C increasing from the point the children were being separated from their parents. Compliance with induction was rated good (lower than 4 points) in 83% of cases in group P, but just 50% of cases in group C. Overall quality was rated much higher (9.3 out of 10) in group P compared with group C (6.8). In group P, all parents said they would repeat the same experience, while in group C, most (87%) said

they would have preferred to stay with their child during anaesthesia.

Dr Sánchez concludes: "Parental presence during anaesthesia induction improves quality of anaesthesia both for parents and children."

She adds: "The idea for this trial started when my colleague was just about to anaesthetise one child who was very upset and crying, hugging his mother on the entrance of the operating room and wanting her to go with him. We thought, 'why not?' and allowed the mother to accompany her child. We did an inhalational anaesthesia procedure for the child while he was still in his mother's arms, and the experience was great. That was the point when we decided to create this trial."

Provided by European Society of Anaesthesiology

Citation: New research shows parental presence improves the quality of child anesthesia (2014, June 1) retrieved 14 July 2023 from

<https://medicalxpress.com/news/2014-06-parental-presence-quality-child-anesthesia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.