

Anal cancer rates quadrupled since mid 70s

5 June 2014



lifestyles has suggested more heterosexual couples are having anal intercourse, which might account for the higher rates of the disease in women. But HPV is a very common virus and any [sexual activity](#) can increase the risk of passing on the infection to a partner.

Great Britain – All ages			
	1975-1977	2009-2011	Percentage change
Male	0.4	1.2	202%
Female	0.4	1.8	374%
Persons	0.4	1.5	299%

(Medical Xpress)—Anal cancer rates in the UK have [increased by nearly 300 per cent](#) over the last 40 years, according to new figures published by Cancer Research UK.

And the increase in cases has been dramatically higher in women (374 per cent) than in men (202 per cent) over this period, rising from 0.4 to 1.8 per 100,000 in females compared to 0.4 to 1.2 per 100,000 in males.

Experts believe the reason for the dramatic rise is likely to be caused by the increasing prevalence of the human papillomavirus (HPV), a virus that is usually transmitted through sexual activity. An estimated 90 per cent of anal cancer cases in the UK are linked to HPV infection.

Overall the [rates](#) have risen from 0.4 in every 100,000 in the mid 70s to 1.5 today. Anal cancer is relatively rare, but awareness of the disease is low as it is still considered taboo by many.

The latest figures also show death rates for anal cancer have more than quadrupled since the mid 70s, with around 6 people now dying every week from the disease in the UK.

Previous research into [sexual attitudes](#) and

Smoking may also be a risk factor for the disease and, though smoking rates have dropped since the 1970s, the drop has been smaller in women than in men, and slower in younger age groups where [anal cancer](#) is more common.

Nick Ormiston-Smith, Cancer Research UK's head of statistics, said: "These are very worrying findings and highlight an increase in a cancer that's not often talked about. Around 1,200 people are diagnosed with cancer of the anus every year in the UK, which means it's still a relatively rare disease. But the rise in incidence, particularly in women, is concerning.

"We don't fully understand the reason for the difference between men and women, but we do know HPV and smoking are strong risk factors for the disease."

The introduction of the HPV vaccine for girls has been an important public health measure and is expected to help reduce the rates of HPV-related cancers in the UK in the future. The Joint

Committee on Vaccination and Immunisation (JCVI) is now considering whether to extend the vaccination programme to boys, as rates of HPV-related cancers are also on the increase in men.

Using a condom also reduces the risk of spreading the infection, but it doesn't completely prevent it from being passed on to a partner.

Jessica Kirby, senior health information manager at Cancer Research UK, said: "Anal cancer is closely linked to HPV, and changes in sexual attitudes mean people are increasingly exposed to the virus. We're not suggesting people take a vow of celibacy, but HPV vaccination, using a condom and being a non-smoker can all help to reduce the risk.

"Early diagnosis is also important so report any unusual or persistent changes in your body or bowel habits to your doctor. The most common symptom is bleeding from the anus. The cause is much more likely to be something less serious but it's important to report symptoms to your doctor and get them checked out."

Provided by Cancer Research UK

APA citation: Anal cancer rates quadrupled since mid 70s (2014, June 5) retrieved 27 May 2021 from <https://medicalxpress.com/news/2014-06-anal-cancer-quadrupled-mid-70s.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.