

# Are the new ACC/AHA guidelines for lipids appropriate for people with diabetes?

June 14 2014

---

New guidelines for the treatment of high cholesterol are not appropriate for people with type 2 diabetes – or are they? Two researchers will debate the topic at a symposium to be held on Tuesday, June 17, at the American Diabetes Association's 74th Scientific Sessions.

At issue are the newest guidelines issued jointly by the American College of Cardiology and American Heart Association in November 2013. Previous guidelines for treating people with [high cholesterol](#) were based upon cutpoints for low-density lipoprotein (LDL) [cholesterol levels](#) (the so-called "bad" cholesterol). People whose cholesterol levels hit those cutpoints were asked to initiate dietary changes, and, if that failed to lower levels sufficiently, statin therapy was recommended.

The new guidelines divide patients into four groups for whom statin treatment is recommended: individuals with cardiovascular disease; individuals with LDL cholesterol (LDL-C) levels higher than 190 mg/dL; individuals with [diabetes](#), between the ages of 40-75, with LDL-C levels 70-189 mg/dL; and individuals without diabetes or cardiovascular disease, with LDL-C 70-189 mg/dL, and an estimated risk of cardiovascular disease greater than 7.5 percent. There are no specific target levels for lowering cholesterol, only the recommendation that statins be used to reduce it.

Under the new guidelines, the vast majority of people with [type 2 diabetes](#) would be recommended for statin therapy. But is that the right call?

Robert H. Eckel, MD, Professor of Medicine and Charles A. Boettcher Chair in Atherosclerosis at the University of Colorado, Anschutz Medical Campus, says, "Yes. Almost all people with diabetes should be on a statin. That's what the evidence tells us. You may not like the new guidelines, but these are evidence-based and this is what the evidence says."

Henry Ginsberg, MD, Irving Professor of Medicine and Director of the Irving Institute for Clinical and Translational Research at Columbia University, disagrees. "The guideline committee used the evidence-based construct much too narrowly," he said.

Ginsberg explained that he agreed with the recommendation to examine a person's risk for cardiovascular disease as well as their cholesterol level when determining whether to prescribe a statin. But he did not agree with the removal of target LDL [cholesterol](#) goals once a person is being treated with medication, or with the lack of recommendations for additional or alternative therapies.

"There are people who can't take a high dose of a statin, or who won't respond as well to a high dose," he said. "In that case, you should be adding another medication. The new guidelines try to make it too simple."

Ginsberg said the new guidelines actually ignore evidence that shows the lower a patient's LDL-C level, the better. "How do you get to those lower levels if statins alone aren't enough?" he asked. "You add a non-statin."

But Eckel said that exceptions, which often come up in patients with diabetes, are addressed in the new guidelines by suggesting that physicians make a judgment call. "It should be an educated judgment, though, based on an understanding of the increased risk of heart disease and stroke for people with diabetes."

The American Diabetes Association recommends that people with diabetes keep LDL-C levels at or below 100 mg/dL, or under 70 mg/dL if they also have [cardiovascular disease](#). The Association recommends using high-dose [statin therapy](#) to reach those goals.

Ginsberg said he agrees with the Association's present guidelines, noting that they are based on a wealth of data that includes, but is not restricted to, randomized clinical trials.

Provided by American Diabetes Association

Citation: Are the new ACC/AHA guidelines for lipids appropriate for people with diabetes? (2014, June 14) retrieved 12 February 2023 from <https://medicalxpress.com/news/2014-06-accaha-guidelines-lipids-people-diabetes.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.