

What you eat may affect your body's internal biological clock

10 July 2014



Food not only nourishes the body but also affects its internal biological clock, which regulates the daily rhythm of many aspects of human behavior and biology. Researchers reporting in the Cell Press journal *Cell Reports* provide new insights into how adjusting the clock through dietary manipulation may help patients with various conditions and show that insulin may be involved in resetting the clock.

An internal biological or 'circadian' clock plays an important role in preferred sleep times, times of peak alertness, and the timing of certain physiological processes. The clock enables maximum expression of genes at appropriate times of the day, allowing organisms to adapt to the earth's rotation. "Chronic desynchronization between physiological and environmental rhythms not only decreases physiological performance but also carries a significant risk of diverse disorders such as diabetes, cardiovascular diseases, [sleep disorders](#), and cancer," says Dr. Makoto Akashi, of Yamaguchi University, in Japan.

The circadian clock involves two major pathways. The first, which responds to light, has been well characterized. The second, which responds to

food, is less understood. Through experiments in cells and mice, Dr. Akashi and his colleagues found, using cell culture, that insulin, a pancreatic hormone that is secreted in response to feeding, may be involved in resetting the circadian clock. "Insulin-mediated phase adjustment of the clock in feeding-relevant tissues may enable the synchronization between mealtime and tissue function, leading to effective digestion and absorption," he says. "In short, insulin may help the stomach clock synchronize with mealtime."

The researchers' findings provide valuable information on how to adjust the circadian clock through dietary manipulation. "For example, for jet lag, dinner should be enriched with ingredients promoting [insulin secretion](#), which might lead to a phase advance of the circadian clock, whereas breakfast would be the opposite," says Dr. Akashi. The findings also suggest that clock adjustments through feeding might not work well in individuals with [insulin resistance](#), a characteristic of patients with type 2 diabetes. Also, there may be side effects related to the [circadian clock](#) when treating patients with insulin.

More information: Cell Reports, Sato et al.: "The role of the endocrine system in feeding-induced tissue-specific circadian entrainment."

[www.cell.com/cell-reports/abstract/2211-1247\(14\)00483-5](http://www.cell.com/cell-reports/abstract/S2211-1247(14)00483-5)

Provided by Cell Press

APA citation: What you eat may affect your body's internal biological clock (2014, July 10) retrieved 22 July 2022 from <https://medicalxpress.com/news/2014-07-affect-body-internal-biological-clock.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.