


Hunger games: How the brain 'browns' fat to aid weight loss

9 October 2014



Yale researchers uncover a molecular process in the brain that controls eating by transforming white fat into brown fat. Credit: Yale University

Researchers at Yale School of Medicine have uncovered a molecular process in the brain known to control eating that transforms white fat into brown fat. This process impacts how much energy we burn and how much weight we can lose. The results are published in the Oct. 9 issue of the journal *Cell*.

Obesity is a rising global epidemic. Excess fatty tissue is a major risk factor for type 2 diabetes, cardiovascular disease, hypertension, neurological disorders, and cancer. People become overweight and obese when [energy](#) intake exceeds [energy expenditure](#), and excess calories are stored in the adipose tissues, which are made up of both white and [brown fat](#). While white fat primarily stores energy as triglycerides, brown fat dissipates chemical energy as heat. The more brown fat you have, the more weight you can lose.

It has previously been shown that energy-storing white fat has the capacity to transform into energy-burning "brown-like" fat. In this new study, researchers from the Yale Program in Integrative

Cell Signaling and Neurobiology of Metabolism, demonstrate that neurons controlling hunger and appetite in the brain control the "browning" of white fat.

Lead author Xiaoyong Yang, associate professor of comparative medicine and physiology at Yale School of Medicine, conducted the study with Tamas Horvath, professor and chair of [comparative medicine](#), and professor of neurobiology and Obstetrics/gynecology at Yale School of Medicine, and their co-authors.

The team stimulated this browning process from the brain in mice and found that it protected the animals from becoming obese on a high-fat diet. The team then studied the molecular changes in hunger-promoting neurons in the hypothalamus and found that the attachment of a unique sugar called "O-GlcNAc" to potassium ion channels acts as a switch to control brain activity to burn fat.

"Our studies reveal white fat "browning" as a highly dynamic physiological process that the brain controls," said Yang. "This work indicates that behavioral modifications promoted by the brain could influence how the amount of food we eat and store in fat is burned."

Yang said hunger and [cold exposure](#) are two life-history variables during the development and evolution of mammals. "We observed that food deprivation dominates over cold exposure in neural control of [white fat](#) browning. This regulatory system may be evolutionarily important as it can reduce heat production to maintain energy balance when we are hungry. Modulating this [brain](#)-to-fat connection represents a potential novel strategy to combat obesity and associated illnesses."

Provided by Yale University

APA citation: Hunger games: How the brain 'browns' fat to aid weight loss (2014, October 9) retrieved 3 May 2021 from <https://medicalxpress.com/news/2014-10-hunger-games-brain-browns-fat.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.