

Transcutaneous electrical nerve stimulation eases vestibulodynia

20 November 2014

(HealthDay)—Transcutaneous electrical nerve stimulation (TENS) may be useful as an additional treatment in the management of therapy-resistant provoked vestibulodynia (PVD), according to research published online Nov. 12 in the *Journal of Sexual Medicine*.

Marleen S. Vallinga, M.D., of the University of Groningen in the Netherlands, and colleagues conducted a longitudinal prospective study of 39 [women](#) (mean age, 27 ± 5.6 years; range: 19 to 41 years) with therapy-resistant PVD. Self-report questionnaires and visual analog scales (VASs) were administered at baseline, following TENS therapy, and at follow-up.

The researchers found that directly following TENS therapy and at follow-up, VAS scores for vulvar pain were significantly reduced from baseline (P

"The addition of self-administered TENS to multidimensional treatment significantly reduced the level of vulvar pain and the need for vestibulectomy," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2014 [HealthDay](#). All rights reserved.

APA citation: Transcutaneous electrical nerve stimulation eases vestibulodynia (2014, November 20) retrieved 8 November 2022 from <https://medicalxpress.com/news/2014-11-transcutaneous-electrical-nerve-eases-vestibulodynia.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.