

Some people may be genetically susceptible to UV tanning dependence

1 December 2014

Researchers have found a possible underlying genetic susceptibility to being dependent on UV tanning.

After interviewing young people about their indoor and outdoor tanning history and using [questionnaires](#) to classify people as being dependent on UV tanning or not, the investigators conducted a large scale scan of approximately 319,000 rare and common genetic variants in the participants' genomes.

"We observed that inherited variation in one gene - known as patched domain containing 2 (PTCHD2) - was significantly associated with whether or not [young people](#), all of whom had previously sunbathed or indoor tanned, exhibited symptoms of tanning dependence," said Dr. Brenda Cartmel, lead author of the *Experimental Dermatology* study.

More information: Cartmel, B., Dewan, A., Ferrucci, L. M., Gelernter, J., Stapleton, J., Leffell, D. J., Mayne, S. T. and Bale, A. E. (2014), Novel gene identified in an exome-wide association study of tanning dependence. *Experimental Dermatology*, 23: 757-759. [DOI: 10.1111/exd.12503](#)

Provided by Wiley

APA citation: Some people may be genetically susceptible to UV tanning dependence (2014, December 1) retrieved 28 July 2022 from <https://medicalxpress.com/news/2014-12-people-genetically-susceptible-uv-tanning.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.