

Non-chest pain presentation doesn't worsen MI outcomes

December 30 2014



(HealthDay)—Among patients with acute ST-elevation myocardial infarction (STEMI), presentation with non-chest pain delays door-to-balloon (DTB) time but does not worsen clinical outcomes, according to research published in the Dec. 15 issue of *The American Journal of Cardiology*.

Jin Sup Park, M.D., of Pusan National University Hospital in Busan, South Korea, and colleagues conducted a prospective observational study of matched cohorts, undergoing primary [percutaneous coronary intervention](#) for STEMI, who presented with chest pain or non-chest pain complaints (976 [patients](#) each). The effect of non-chest pain complaints on DTB time and clinical outcomes was assessed.

The researchers found that patients with non-chest pain complaints, compared with those with chest pain, experienced significant delays in

DTB time (median, 84 versus 74 minutes, respectively; P

"In conclusion, short- and long-term [clinical outcomes](#) in patients with STEMI with non-chest pain complaints do not differ from those of patients with [chest pain](#) as the presenting symptom, despite having delayed diagnosis and reperfusion," the authors write.

More information: [Abstract](#)

[Full Text \(subscription or payment may be required\)](#)

Copyright © 2014 [HealthDay](#). All rights reserved.

Citation: Non-chest pain presentation doesn't worsen MI outcomes (2014, December 30)
retrieved 26 April 2023 from
<https://medicalxpress.com/news/2014-12-non-chest-pain-doesnt-worsen-mi.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--