

Aerobic exercise reduces fatigue with radiotherapy for breast CA

15 January 2015



More information: [Abstract](#)
[Full Text](#)

Copyright © 2014 [HealthDay](#). All rights reserved.

(HealthDay)—An aerobic exercise program can reduce fatigue in women undergoing radiotherapy for breast cancer, according to a study published in the January issue of the *Journal of Clinical Nursing*.

Tsui-Yun Yang, R.N., from the National Taipei University in Taiwan, and colleagues assigned 28 consecutive, willing participants with early-stage [breast cancer](#) to an intervention of six weeks of mild- to moderate-intensity aerobic exercise training during their [radiotherapy](#). The next 30 consecutive participants were assigned to routine care without special exercise.

The researchers observed no group differences in baseline fatigue severity, fatigue interference, or hemoglobin levels. There was a significant group-by-time-interaction effect noted for fatigue severity and interference ($P < 0.001$), with the fatigue severity and interference decreasing significantly over time for women in the intervention group but increasing over time for women in the control group.

"The mild- to moderate-intensity [aerobic exercise](#) program reduces the fatigue of Taiwanese women with breast cancer undergoing radiotherapy," the authors write.

APA citation: Aerobic exercise reduces fatigue with radiotherapy for breast CA (2015, January 15)
retrieved 2 July 2022 from <https://medicalxpress.com/news/2015-01-aerobic-fatigue-radiotherapy-breast-ca.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.