

Are antipsychotic drugs more dangerous to dementia patients than we think?

18 March 2015

Drugs aimed at quelling the behavior problems of dementia patients may also hasten their deaths more than previously realized, a new study finds.

The research adds more troubling evidence to the case against [antipsychotic drugs](#) as a treatment for the delusions, hallucinations, agitation and aggression that many people with Alzheimer's disease and other dementias experience.

In the new issue of the journal *JAMA Psychiatry*, researchers report findings from nearly 91,000 American veterans over the age of 65 with dementia.

Data from each patient who took a psychiatric medication was matched to data from a similar patient who didn't take drugs to calm behavior, by researchers from the University of Michigan Medical School and VA Center for Clinical Management Research.

Those taking drugs called [antipsychotics](#) had outsize risks of death. Among those taking the newer, more commonly used antipsychotics, the risk climbed along with the dose.

The study also examined other [psychiatric medications](#). The risk of death seen with the mood stabilizer [valproic acid](#) was similar to the antipsychotics. Antidepressants had less risk compared with antipsychotics and valproic acid, but it was still higher than that of those not taking any psychiatric medications to treat behavior issues in dementia.

Antipsychotic drugs have significant risk of side effects, and the U.S. Food and Drug Administration warns that their use in people with dementia is associated with increased risk for cardiovascular adverse events and the risk of death.

"The harms associated with using these drugs in dementia patients are clear, yet clinicians continue

to use them," says lead author and U-M/VA psychiatrist Donovan Maust, M.D., M.S. "That's likely because the symptoms are so distressing. These results should raise the threshold for prescribing further."

This new study and a recent state-of-the-art review by Maust's co-author Helen Kales, M.D., show that mortality risk of antipsychotics is higher, and benefit lower, than previously thought.

But the distress caused by uncontrolled symptoms probably explains why one-third of older adults with dementia who had long-term nursing home stays in 2012 were prescribed an [antipsychotic medication](#), according to a recent U.S. Government Accountability Office report. About 14 percent of [dementia patients](#) in the community were prescribed an antipsychotic that same year, the GAO found.

Patients, families and their care teams deserve better to prevent and treat distressing dementia symptoms, Maust says.

The "DICE" approach to assess and manage behavioral symptoms in dementia that has been put forth by Kales and colleagues could help.

This approach emphasizes putting non-pharmacological strategies first. But the approach takes more time than writing a prescription, and its use will depend on the support of policy-makers and alignment of reimbursement strategies.

"In other words, non-pharmacologic approaches will only succeed if we as a society agree to pay front-line providers for the time needed to 'do the right thing'," says Kales.

Maust, Kales and their colleagues hope to continue studying the issue of [mortality risk](#) from the use of psychotropic drugs in dementia, and prescription patterns for these drugs, using other datasets.

More information: *JAMA Psychiatry*. 2015;72(5).

[DOI: 10.1001/jamapsychiatry.2014.3018](https://doi.org/10.1001/jamapsychiatry.2014.3018)

Provided by University of Michigan Health System

APA citation: Are antipsychotic drugs more dangerous to dementia patients than we think? (2015, March 18) retrieved 12 October 2022 from <https://medicalxpress.com/news/2015-03-antipsychotic-drugs-dangerous-dementia-patients.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.