

# Can a parent's concerns predict autism?

23 April 2015



Lori Sacrey and Lonnie Zwaigenbaum. Credit: Faculty of Medicine & Dentistry, University of Alberta

As co-director of the University of Alberta's Autism Research Centre, Lonnie Zwaigenbaum has devoted much of his career to understanding how to identify autism as early as possible. But despite his years of experience, Zwaigenbaum says many physicians like him would do well to seek other expert advice when working with children not yet diagnosed—that of the parents of these young patients.

"Parents are the experts when it comes to their kids and their [observations](#) are really valuable," says Zwaigenbaum. "In some respects, parents are picking up on differences at six and nine months of age that we have a much harder time seeing in the clinic."

Zwaigenbaum, a professor in the Faculty of Medicine & Dentistry's Department of Pediatrics, along with Lori Sacrey, a postdoctoral fellow in the Department of Pediatrics, are the authors of a new study published in the March edition of the *Journal of the American Academy of Child & Adolescent Psychiatry*. The study found parents of children at a high risk for [autism spectrum disorder](#) (ASD) not only reported more early concerns than other parents, but also that those concerns can be predictive of autism spectrum disorder.

The researchers examined the reported concerns of parents from about 300 families for children aged six months to three years old over a 12-year period. The sample included the concerns of parents of children without any heightened risk of ASD, as well as the parents of children at high risk (who had an older sibling already diagnosed with ASD). At three years of age all of the children underwent a clinical assessment to determine if they had autism or not. Researchers then looked back at the parent concerns to see if the groups differed on the types of concerns parents had, as well as the number of concerns they had.

"We found that parents whose children ended up being diagnosed at three years of age did report more concerns," says Sacrey. "Interestingly, they reported sensory and motor concerns starting at the age of six months. And then they increasingly reported more language and social concerns at about 12-15 months of age."

"It really highlights the importance of talking to [parents](#) and taking their concerns seriously," adds Zwaigenbaum.

The researchers believe that by acting on parental concerns early, health professionals can provide better care for [children](#) at risk of autism spectrum disorder.

"Where interventions are concerned, the earlier you can start with the patients, the better the prognosis is," explains Sacrey. "If you can identify a child at a heightened risk earlier, before their first birthday, then you can start working with them to address early developmental difficulties, which can ultimately enhance their skill development and improve their outcomes."

"Parents play a critical role in implementing these interventions, building learning opportunities into everyday caregiving and play activities."

Provided by University of Alberta Faculty of

Medicine & Dentistry

APA citation: Can a parent's concerns predict autism? (2015, April 23) retrieved 24 June 2022 from <https://medicalxpress.com/news/2015-04-parent-autism.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*