

# RYGB cuts markers of oxidative stress in subcutaneous tissue

29 May 2015



plasma," the authors write. "The results of the present study indicate that it has a similar effect in subcutaneous adipose tissue."

**More information:** [Abstract](#)  
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

(HealthDay)—Roux-en-Y gastric bypass (RYGB) surgery attenuates markers of oxidative stress in subcutaneous adipose tissue, according to a study published online May 22 in *Diabetes*.

X. Julia Xu, Ph.D., from the Boston University School of Medicine, and colleagues examined how AMP-activated protein kinase (AMPK) activity, oxidative stress, and inflammation are affected by RYGB surgery. Data were included from 11 patients who were examined immediately prior to and three months after surgery.

The researchers found that AMPK activity increased 3.5-fold after surgery, and oxidative stress declined by 50 percent in [subcutaneous adipose tissue](#). There was an 80 percent reduction in malonyl-coA levels. Patients also had improvements in their [body mass index](#), insulin sensitivity (measured by the homeostasis model assessment), and increased circulating high molecular weight adiponectin; there was also a reduction in fasting plasma insulin levels. Postoperatively, there was no change in the expression of inflammatory markers in subcutaneous tissue, but plasma C-reactive protein was reduced by 50 percent.

"Bariatric surgery has been demonstrated to attenuate markers of [oxidative stress](#) in liver and

APA citation: RYGB cuts markers of oxidative stress in subcutaneous tissue (2015, May 29) retrieved 22 July 2022 from

<https://medicalxpress.com/news/2015-05-rygb-markers-oxidative-stress-subcutaneous.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*