

Early healthy eating intervention in pregnancy helps obese women

16 July 2015



early HE interventions in obese [pregnant women](#)," the authors write.

One author disclosed financial ties to Novo Nordisk.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

(HealthDay)—A healthy eating (HE) intervention is associated with lower gestational weight gain (GWG) and fasting glucose than a physical activity (PA) intervention, according to a study published online June 25 in *Diabetes Care*.

In a multicenter randomized trial, David Simmons, M.D., from Addenbrookes Hospital in Cambridge, U.K., and colleagues compared the impact of three lifestyle interventions on [gestational diabetes mellitus](#) (GDM) risk. One hundred fifty pregnant women ([body mass index](#), ≥ 29 kg/m²) without GDM were randomized to HE, PA, or HE + PA. Women received five face-to-face and four optional telephone-coaching sessions.

The researchers found that 32 percent of the trial participants developed GDM by 35 to 37 weeks, and 20 percent achieved GWG of less than 5 kg. Compared with those in the PA group, women in the HE group had less GWG (-2.6 kg; $P = 0.03$) and lower fasting glucose (-0.3 mmol/L; $P = 0.01$) at 24 to 28 weeks. Differences were not significant for the HE + PA group versus other groups.

"Although a larger trial is still clearly needed, these pilot findings are promising and support the use of

APA citation: Early healthy eating intervention in pregnancy helps obese women (2015, July 16) retrieved 30 July 2022 from <https://medicalxpress.com/news/2015-07-early-healthy-intervention-pregnancy-obese.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.