

Many with nonceliac wheat sensitivity have autoimmune diseases

August 27 2015



(HealthDay)—More patients with nonceliac wheat sensitivity (NCWS) and celiac disease (CD) than irritable bowel syndrome (IBS) develop autoimmune diseases (ADs), according to a study published in the September issue of *Gastroenterology*.

Antonio Carroccio, M.D., from the University of Palermo in Italy, and colleagues conducted a retrospective study of 131 [patients](#) with NCWS and 151 control patients with CD and IBS to examine the prevalence of AD and antinuclear antibodies (ANA). A prospective study was conducted involving 42 patients with NCWS and 100 age- and sex-matched controls with CD or IBS.

The researchers found that 29 percent of patients with NCWS and CD developed ADs (mainly Hashimoto's thyroiditis) compared to 4 percent of patients with IBS (P retrospective study. The corresponding values in

the prospective study were 28, 7.5, and 6 percent ($P = 0.02$ for CD and $P = 0.005$ for IBS versus NCWS).

"Higher proportions of patients with NCWS or CD develop autoimmune disorders, are ANA positive, and showed DQ2/DQ8 haplotypes compared with patients with IBS," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

Citation: Many with nonceliac wheat sensitivity have autoimmune diseases (2015, August 27) retrieved 15 July 2023 from <https://medicalxpress.com/news/2015-08-nonceliac-wheat-sensitivity-autoimmune-diseases.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.