

Allergy to soy reported after percutaneous sensitization

September 15 2015



(HealthDay)—Reduced skin barrier function seen in patients with atopic dermatitis may cause percutaneous sensitization by various proteins, such as food, according to a case report published in the September issue of *The Journal of Dermatology*.

Akiko Yagami, M.D., from Fujita Health University School of Medicine in Japan, and colleagues described the case of a 30-year-old esthetician with atopic dermatitis who developed anaphylactic symptoms after eating soy. She had no history of hand eczema or soy food allergy before working as an esthetician; she started experiencing symptoms of itchy eczema on her fingers several months after starting to touch cosmetic lotions at age 23 years. She experienced severe symptoms including urticaria and dyspnea after eating soy products at age 28 years.

The researchers found that in laboratory tests, she had specific



immunoglobulin E antibodies for soy, Japanese cedar, and Japanese white birch. A skin prick test was positive for soy extract, the cosmetic lotion used by the patient, and a commercially available soy milk. In three healthy controls, skin prick tests of the same cosmetic lotion yielded negative results.

"The patient with atopic dermatitis, which tends to reduce the barrier function of the <u>skin</u>, was percutaneously sensitized to soy protein through frequent contact with cosmetic lotions containing soy-based ingredients in a humid environment while working as an esthetician and she subsequently developed anaphylactic reaction to <u>soy</u>," the authors write.

More information: Abstract

Full Text (subscription or payment may be required)

Copyright © 2015 HealthDay. All rights reserved.

Citation: Allergy to soy reported after percutaneous sensitization (2015, September 15) retrieved 20 February 2023 from

https://medicalxpress.com/news/2015-09-allergy-soy-percutaneous-sensitization.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.