

ASTRO: Shorter radiation OK for early breast cancers

21 October 2015



cancers."

More information: [Abstract](#)
[More Information](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

(HealthDay)—Hypofractionated radiation therapy with incorporated boost over a shorter period of time treats early-stage breast cancer as well as longer, conventional radiation therapy, a new study suggests. The findings were scheduled to be presented at the annual meeting of the American Society for Radiation Oncology, held from Oct. 18 to 21 in San Antonio.

Researchers from Fox Chase Cancer Center-Temple Health in Philadelphia report that eight years after treatment, there were no significant differences in survival among hundreds of patients who received hypofractionated whole-breast radiation with incorporated boost over four weeks or conventional fractionation with sequential boost.

"This is good news for cancer patients because a shorter length of treatment is not only more convenient, but it helps to enable a woman to get her life back to normal after [breast cancer](#) treatment," Stephanie Bernik, M.D., chief of surgical oncology at Lenox Hill Hospital in New York City, told *HealthDay*.

"At this time, the shorter [radiation therapy](#) is offered to a select group of women," Bernik added. "With more and more studies showing equivalence, the hope is that this therapy can be expanded to women with even more aggressive

APA citation: ASTRO: Shorter radiation OK for early breast cancers (2015, October 21) retrieved 4 May 2021 from <https://medicalxpress.com/news/2015-10-astro-shorter-early-breast-cancers.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.