

Sexual dysfunction is prevalent among recently deployed veterans

November 4 2015

In a recent study of 247 US veterans returning from conflicts in Iraq and Afghanistan, almost 18% screened positive for sexual functioning difficulties. Self-reported sexual dysfunction was most strongly linked with depression, post-traumatic stress disorder, and female sex.

Additional research should examine the causes of specific sexual health concerns in returning veterans, which might relate to dysfunction due to injury or events prior to, during, or following military service.

"Sexual dysfunction is as prevalent as other conditions we think of as being related to military service, such as post-traumatic stress disorder and traumatic brain injury," said Dr. Drew Helmer, co-author of the *Sexual Medicine* study. "Recent combat veterans are often at a stage of life when sexual health is critical for establishing and sustaining relationships and starting families. We should make it easy for all veterans to discuss sexual health concerns with their healthcare providers and receive effective care."

More information: Gregory R. Beaulieu et al. An Exploration of Returning Veterans' Sexual Health Issues Using a Brief Self-Report Measure, *Sexual Medicine* (2015). [DOI: 10.1002/sm2.92](https://doi.org/10.1002/sm2.92)

Provided by Wiley

Citation: Sexual dysfunction is prevalent among recently deployed veterans (2015, November 4)
retrieved 1 February 2023 from
<https://medicalxpress.com/news/2015-11-sexual-dysfunction-prevalent-deployed-veterans.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.