

Sleep-disordered breathing in REM linked to insulin resistance

November 5 2015



(HealthDay)—The correlation between sleep-disordered breathing (SDB) and glucose metabolism varies for rapid eye movement (REM) sleep and non-REM sleep, according to a study published in the Nov. 1 issue of the *American Journal of Respiratory and Critical Care Medicine*.

Hassan A. Chami, M.D., from the American University of Beirut, and colleagues characterized the association between REM-related SDB, glucose intolerance, and insulin resistance in a community-based sample of 3,310 participants. The apnea-hypopnea index (AHI) was used to quantify SDB severity during REM (AHI_{REM}) and non-REM (AHI_{NREM}) sleep. A glucose tolerance test assessed fasting and two-hour post-challenge glucose levels in 2,264 participants; the homeostatic model assessment index for insulin resistance (HOMA-IR) was measured for 1,543 participants.



The researchers found that in models that adjusted for age, sex, race, and site, AHI_{REM} and AHI_{NREM} correlated with fasting glycemia, postprandial glucose levels, and HOMA-IR. AHI_{REM} was only associated with HOMA-IR and AHI_{NREM} was only associated with fasting and postprandial glucose levels after further adjustment for body mass index, waist circumference, and sleep duration.

"AHI_{REM} is associated with <u>insulin resistance</u> but not with fasting glycemia or glucose intolerance," the authors write.

More information: <u>Full Text (subscription or payment may be required)</u>

Copyright © 2015 HealthDay. All rights reserved.

Citation: Sleep-disordered breathing in REM linked to insulin resistance (2015, November 5) retrieved 15 July 2023 from https://medicalxpress.com/news/2015-11-sleep-disordered-rem-linked-insulin-resistance.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.