

No direct survival effect for moderate drinking in seniors

25 November 2015



direct protective effect on mortality," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2015 [HealthDay](#). All rights reserved.

(HealthDay)—For older adults, light-to-moderate drinking does not have a protective effect on mortality after adjustment for health status and physical activity, according to a study published online Nov. 23 in the *Journal of the American Geriatrics Society*.

Antonio Muscari, M.D., from the University of Bologna in Italy, and colleagues examined the correlation between light-to-moderate alcohol consumption and [mortality](#) in a prospective, longitudinal study among individuals aged 65 and older. Data were included for 2,318 abstainers and 2,309 light-to-moderate drinkers (no more than two alcoholic units/day). Follow-up information was obtained from 2,752 survivors six years later; mortality information was obtained from death certificates.

The researchers observed independent correlations for male sex, being physically active, and good health status with light-to-moderate drinking (P protective effect of light-to-moderate drinking on mortality; this persisted after adjustment for age, sex, risk factors, and cardiovascular events (adjusted hazard ratio, 0.77; 95 percent confidence interval, 0.68 to 0.88; P

"After accounting for [health status](#) and [physical activity](#), light-to-moderate alcohol drinking had no

APA citation: No direct survival effect for moderate drinking in seniors (2015, November 25) retrieved 3 May 2021 from <https://medicalxpress.com/news/2015-11-survival-effect-moderate-seniors.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.