

Paracentesis underutilized in patients with cirrhosis, ascites

11 December 2015

Copyright © 2015 [HealthDay](#). All rights reserved.



(HealthDay)—For patients admitted with diagnoses of cirrhosis and ascites, paracentesis is associated with decreased in-hospital mortality but is underutilized, according to a study published online Dec. 8 in the *Journal of Gastroenterology and Hepatology*.

John N. Gaetano, M.D., from the University of Chicago, and colleagues identified 31,614 adults, non-electively admitted with diagnoses of cirrhosis and ascites. Utilization and outcomes of paracentesis were examined.

The researchers found that 51 percent of the hospitalized patients underwent paracentesis. The overall in-hospital [mortality](#) rate was 7.6 percent, with increased mortality seen among patients who did not versus those who did undergo paracentesis (8.9 versus 6.3 percent; P patients diagnosed with spontaneous bacterial peritonitis (both P

"[These data support] the use of paracentesis as a key inpatient quality measure among hospitalized adults with [cirrhosis](#)," the authors write. "Future studies are needed to investigate the barriers to paracentesis use on admission."

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

APA citation: Paracentesis underutilized in patients with cirrhosis, ascites (2015, December 11) retrieved 18 June 2021 from <https://medicalxpress.com/news/2015-12-paracentesis-underutilized-patients-cirrhosis-ascites.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.