

Trends in insulin use, glycemic control explored

8 January 2016



for those on oral medication only (7.0 to 8.5 years; $P = 0.36$). Among patients with any insulin use, 31.4 percent had an hemoglobin A1c (HbA1c)

(HealthDay)—Since 1988, the proportion of patients with diabetes currently on any insulin has remained stable, according to research published online Dec. 30 in *Diabetes Care*.

Elizabeth Selvin, Ph.D., from the Johns Hopkins Bloomberg School of Public Health in Baltimore, and colleagues examined national trends in glycemic control by [insulin](#) and oral diabetes medication use. Data were included for 4,947 adult participants with diagnosed diabetes in the National Health and Nutrition Examination Survey for the 1988 to 1994 and 1999 to 2012 cycles.

The researchers found that the proportion of patients with diabetes currently on insulin was relatively stable (29.1 percent in 2005 to 2012; 30.3 percent in 1988 to 1994; and 24.8 percent in 1999 to 2004). Diabetes duration was a strong independent determinant of insulin use. During the study period, the duration of diabetes increased significantly for those on any insulin (12.7 to 17.3 years; $P < 0.001$), compared with diabetes duration

APA citation: Trends in insulin use, glycemic control explored (2016, January 8) retrieved 28 April 2021 from <https://medicalxpress.com/news/2016-01-trends-insulin-glycemic-explored.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.