

Stress could help activate brown fat

9 February 2016



Credit: George Hodan/public domain

Mild stress stimulates the activity and heat production by brown fat associated with raised cortisol, according to a study published today in *Experimental Physiology*.

Brown adipose tissue (BAT), also known as [brown fat](#), is one of two types of fat found in humans and other mammals. Initially only attributed to babies and hibernating mammals, it was discovered in recent years that adults can have brown fat too. Its main function is to generate body heat by burning calories (opposed to [white fat](#), which is a result of storing excess calories. People with a lower [body mass index](#) (BMI) therefore have a higher amount of brown fat.

To induce a mild psychological [stress](#), five healthy lean women had to solve a short maths test in the first run, but in the second run, the test was substituted with a relaxation video. To assess stress responses, the scientists measured cortisol in the saliva. To measure the activity of brown fat, the researchers used infrared thermography to detect changes in temperature of the skin overlying the main area of brown fat in humans (in the neck (supraclavicular) region).

Although the actual maths tests did not elicit an acute stress response, the anticipation of being tested did, and led to raised cortisol and warmer brown fat. Both were positively correlated, with higher cortisol linked with more fat activity and thus more potential [heat production](#).

Prof Michael E Symonds from The School of Medicine, University of Nottingham and Co-author of the study comments,

'Our research indicates that the variation in brown fat activity between individuals may be explained by differences in their response to psychological stress. This is important as brown fat has a unique capacity to rapidly generate heat and metabolise glucose.

'Most adults only have between 50-100 g of brown fat but because its capacity to generate heat is 300 times greater (per unit mass) than any other tissue, brown fat has the potential to rapidly metabolise glucose and lipids. There is an inverse relationship between the amount of brown fat and BMI, and whether this is a direct consequence of having more active fat remains to be fully established.

'A better understanding of the main factors controlling brown fat activity, which include diet and activity, therefore has the potential to introduce sustainable interventions designed to prevent obesity and diabetes. In future, new techniques to induce [mild stress](#) to promote brown fat activity could be incorporated alongside dietary and/or environmental interventions. This is likely to contrast with the negative effects of chronic and more severe stress that can contribute to poor metabolic health.'

More information: Lindsay J Robinson et al. Brown adipose tissue activation as measured by infrared thermography by mild anticipatory psychological stress in lean healthy females, *Experimental Physiology* (2016). DOI: [10.1113/EP085642](https://doi.org/10.1113/EP085642)

Provided by The Physiological Society

APA citation: Stress could help activate brown fat (2016, February 9) retrieved 16 August 2022 from <https://medicalxpress.com/news/2016-02-stress-brown-fat.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.