

Australian hoarders falling between the gaps

February 9 2016



A leading expert on hoarding disorder called for a major rethink in the way hoarding is managed in Australia. Credit: Thomas Vandenberghe, Flickr

A leading expert on hoarding disorder has developed a new treatment for the condition and called for a major rethink in the way hoarding is managed in Australia.

Professor Mike Kyrios, Director of The Australian National University (ANU) Research School of Psychology, said people with hoarding

disorder are falling between the gaps of Australia's medical system.

Hoarding disorder affects between two to six per cent of the population. People with the condition excessively acquire or save items and have difficulty getting rid of possessions regardless of the value. This then leads to clutter in the home or [work spaces](#) rendering those spaces unusable and potentially dangerous.

"This is a severe mental health condition that affects people's lives and is associated with unnecessary deaths. We know that 25 per cent of deaths from fires in homes occur in the homes of people with a hoarding problem," Professor Kyrios said.

"Existing Medicare services only allow for 10 sessions of treatment, where hoarding takes a much longer time to treat - anywhere between six-months and twelve-months.

"Effective treatments do exist however there are very few people in Australia who can provide them.

"We need a much more multi-disciplinary approach to the management of hoarding in the community and we need to teach a range of professionals how to manage it effectively."

Once considered to be a symptom of Obsessive Compulsive Disorder (OCD), in 2013 'hoarding disorder' was added to the American Psychiatric Association's Diagnostic and Statistical Manual of Mental Disorders.

Professor Kyrios and his team have played a lead role in the development of a psychological treatment of hoarding for use in Australia, outlined in a paper published in the latest Journal of Clinical Psychology & Psychotherapy, and is currently trialling online treatment

supports.

He said older Australians were particularly susceptible to the effects of [hoarding disorder](#) and that there were a number of aspects that lead to people developing the condition including genetic and experiential factors.

"In many instances people who feel insecure about themselves as a result of their early upbringing, will overly acquire things, to defend themselves against the threats they feel are out there in the world," Professor Kyrios said.

"It's almost as if they build a defensive wall around themselves through the use of objects. However, we can teach people to have a more positive relationship with possessions and to minimise the clutter in their homes."

More information: Richard Moulding et al. Short-Term Cognitive-Behavioural Group Treatment for Hoarding Disorder: A Naturalistic Treatment Outcome Study, *Clinical Psychology & Psychotherapy* (2016). [DOI: 10.1002/cpp.2001](https://doi.org/10.1002/cpp.2001)

Provided by Australian National University

Citation: Australian hoarders falling between the gaps (2016, February 9) retrieved 10 January 2023 from <https://medicalxpress.com/news/2016-02-australian-hoarders-falling-gaps.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--