

# Cox2 inhibition improves preeclampsia symptoms in a mouse model

17 March 2016

---

Preeclampsia is characterized by elevated blood pressure in the second half of pregnancy and impaired blood flow to the placenta, which increases the risk of premature birth and pregnancy complications.

In this issue of *JCI Insight*, Robin Davisson and colleagues use a mouse model that spontaneously develops cardinal features of preeclampsia to explore its underlying causes. They show that even before preeclamptic symptoms develop, embryos have implantation defects that are associated with increased levels of the pro-inflammatory molecule cyclooxygenase 2 (Cox2).

Treatment of mice with the Cox2 inhibitor celecoxib prior to implantation resulted in more typical implantation features, improved fetal growth, and reduced gestational hypertension.

These findings support further exploration of Cox2 inhibition early in pregnancy as an approach to prevent [preeclampsia](#).

**More information:** Jenny L. Sones et al. Decidual Cox2 inhibition improves fetal and maternal outcomes in a preeclampsia-like mouse model, *JCI Insight* (2016). DOI: [10.1172/jci.insight.75351](https://doi.org/10.1172/jci.insight.75351)

Provided by Journal of Clinical Investigation

APA citation: Cox2 inhibition improves preeclampsia symptoms in a mouse model (2016, March 17) retrieved 26 July 2022 from <https://medicalxpress.com/news/2016-03-cox2-inhibition-preeclampsia-symptoms-mouse.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*