

# Central corneal thickness influenced by body position

March 21 2016

---



(HealthDay)—Central corneal thickness (CCT) is influenced by body

position, with a decrease noted in the first 30 minutes of supine positioning, according to a study published online March 14 in *Clinical & Experimental Ophthalmology*.

Jessica S. Maslin, M.D., from the Yale University School of Medicine in New Haven, Conn., and colleagues conducted a cross-sectional study involving 23 [patients](#) with open-angle [glaucoma](#) and 23 healthy subjects. The authors measured CCT using an ultrasound pachymeter in each subject. Three consecutive measurements were taken in each eye in the sitting position, and in a supine position after 10 and 30 minutes.

The researchers found that CCT decreased with supine positioning at 10 and 30 minutes in healthy subjects (mean =  $-5.2 \mu\text{m}$  [P = 0.0043] and  $-6.5 \mu\text{m}$  [P "CCT is a dynamic measurement that can be influenced by [body position](#)," the authors write. "It decreases linearly in the first 30 minutes of supine positioning at a similar rate in both open-angle glaucoma patients and in healthy subjects."

**More information:** [Abstract](#)  
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

Citation: Central corneal thickness influenced by body position (2016, March 21) retrieved 13 February 2023 from <https://medicalxpress.com/news/2016-03-central-corneal-thickness-body-position.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--