

Helping people with Parkinson's get a better night's sleep

April 11 2016, by David Stacey



A volunteer with the West Australian Participant Pool (WAPP) who does not have Parkinson's, Ian Murray, demonstrates some of the equipment used in Parkinson's sleep studies. Mr Murray is wearing a home-based polysomnography device which measures sleep and breathing.

A new study at The University of Western Australia may offer hope in alleviating some of the memory and thinking skills problems associated with the debilitating movement disorder, Parkinson's.

It comes as the Shake It Up Australia Foundation urges Australians to support World Parkinson's Day on April 11 to help raise awareness for those living with Parkinson's, along with money for vital research.

UWA PhD student Maria Pushpanathan said the research, which

examined the links between poor [sleep](#) and cognition in people with Parkinson's and was published in *Sleep Medicine Reviews*, found that disturbed sleep had a profound effect on a patient's concentration, memory and planning ability.

The research was undertaken as part of the ParkC project, a longitudinal study of thinking and motor symptoms in Parkinson's which is a collaboration between Curtin University and UWA.

Ms Pushpanathan and co-authors, Associate Professor Romola Bucks and ParkC director Dr Andrea Loftus, carried out a 'meta-analysis', or a statistical examination, of all the available data from independent studies on night-time sleep problems and cognition in people with Parkinson's.

"While Parkinson's is classified as a movement disorder, it also affects thinking and memory skills, and causes sleep disruption," Ms Pushpanathan said.

"It turns out that these sleep problems could be a significant factor leading to cognitive issues like poorer attention and memory loss. Having Parkinson's is tough, let alone if we add more cognitive problems due to [poor sleep](#)."

Ms Pushpanathan said sleep disorders affect up to 98 per cent of people with Parkinson's with the range of common disturbances including insomnia, [sleep fragmentation](#), sleep-related breathing [disorders](#), restless leg syndrome, REM sleep behaviour disorder and nightmares.

"While the number of studies that fit our criteria was relatively small and this field of research is in its infancy, it is the beginning of a new scientific adventure," Ms. Pushpanathan said.

"What we have found could have major clinical implications by helping

to raise awareness of how problematic sleep can have an enormous impact in Parkinson's."

"Our results also suggest that many types of sleep disorder, such as insomnia, and sleep-related breathing disorder, are not being identified or studied well in Parkinson's.

"We know that sufficient, good quality sleep is vital for physical and mental health, as well as for thinking and memory. Further research investigating the impact of these common [sleep disorders](#) in Parkinson's is desperately needed."

Ms Pushpanathan said researchers now hope to look at how sleep disorder evolves as Parkinson's develops.

"Approximately 70, 000 Australians are living with Parkinson's - imagine how life would be different if sleep was no longer a struggle, and [memory](#) loss could be delayed, reduced, or even prevented," she said.

More information: Maria E. Pushpanathan et al. The relationship between sleep and cognition in Parkinson's disease: A meta-analysis, *Sleep Medicine Reviews* (2016). [DOI: 10.1016/j.smrv.2015.04.003](https://doi.org/10.1016/j.smrv.2015.04.003)

Provided by University of Western Australia

Citation: Helping people with Parkinson's get a better night's sleep (2016, April 11) retrieved 13 July 2023 from <https://medicalxpress.com/news/2016-04-people-parkinson-night.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is

provided for information purposes only.