

# Commonly used reflux and ulcer medication may cause serious kidney damage

14 April 2016

New research indicates that long-term use of certain medications commonly used to treat heartburn, acid reflux, and ulcers can have damaging effects on the kidneys. The findings come from a study appearing in an upcoming issue of the *Journal of the American Society of Nephrology (JASN)*.

In 2013, an estimated 15 million Americans were prescribed [proton pump](#) (PPIs), which reduce gastric acid production. This number is likely an underestimate because the medications are also available over-the-counter and can be purchased without prescription.

To assess the safety of this widely used drug class, a team led by Yan Xie, MPH and Ziyad Al-Aly, MD, FASN (Clinical Epidemiology Center at the VA Saint Louis Health Care System and Washington University in Saint Louis) analyzed information from the Department of Veterans Affairs national databases. The investigators identified 173,321 new users of PPIs and 20,270 new users of histamine H2 [receptor blockers](#), an alternative class of drugs also used to suppress [stomach acid](#). Over 5 years of follow-up, those taking PPIs were more likely to experience kidney function decline than those taking H2 receptor blockers. PPI users also had a 28% increased risk of developing chronic kidney disease and a 96% increased risk of developing kidney failure. Furthermore, there was a graded association between duration of PPI use and risk of kidney problems, with those who took PPIs for a longer time being more likely to develop kidney issues.

The findings suggest that long-term use of PPIs may be harmful to the kidneys and should be avoided. PPI use may not only increase the risk of developing chronic kidney disease, but may also increase the risk of its progression to complete [kidney](#) failure. "The results emphasize the importance of limiting PPI use only when it is medically necessary, and also limiting the duration

of use to the shortest duration possible," said Dr. Al-Aly. "A lot of patients start taking PPIs for a medical condition, and they continue much longer than necessary."

The results also provide insights for future investigations on drug safety. "The study serves as a model to leverage the availability of Big Data—with VA data being a prime example—and advanced analytics to determine long term safety profiles of commonly used medications and promote pharmacovigilance," said Xie.

**More information:** "Proton Pump Inhibitors and Risk of Incident and Progression of Chronic Kidney Disease and ESRD," April 14, 2016, [DOI: 10.1681/ASN.2015121377](#)

Provided by American Society of Nephrology

APA citation: Commonly used reflux and ulcer medication may cause serious kidney damage (2016, April 14) retrieved 9 July 2022 from <https://medicalxpress.com/news/2016-04-commonly-reflux-ulcer-medication-kidney.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*