

# No link for tea, coffee intake with barrett's esophagus

5 May 2016



correlation between the risk of BE and consumption of coffee (adjusted odds ratio, 1.04; 95 percent confidence interval, 0.76 to 1.42) or tea (adjusted odds ratio, 1.11; 95 percent confidence interval, 0.85 to 1.44).

"These data do not support an association between consumption of coffee or tea and the risk of BE," the authors write. "It is unlikely that avoidance of [coffee](#) or tea will protect against BE."

**More information:** [Abstract](#)

[Full Text](#)

[Editorial](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

(HealthDay)—After adjustment for confounding variables there is no correlation between the risk of Barrett's esophagus (BE) and tea or coffee consumption, according to a study published in the May issue of *Clinical Gastroenterology and Hepatology*.

Krishna C. Sajja, M.D., from the Baylor College of Medicine in Houston, and colleagues examined the correlation between tea/[coffee consumption](#) and BE in a cross-sectional study among U.S. veterans. Three hundred ten patients with histologically confirmed BE were compared with 1,728 individuals with no endoscopic or [histopathologic features](#) of BE.

The researchers observed a statistically significant association between the risk of BE with consumption of coffee (odds ratio, 1.41; 95 percent confidence interval, 1.06 to 1.87) or tea (odds ratio, 1.34; 95 percent confidence interval, 1.05 to 1.71) in univariate models. However, after adjustment for confounders, including sex and race, there was no

APA citation: No link for tea, coffee intake with barrett's esophagus (2016, May 5) retrieved 2 December 2022 from <https://medicalxpress.com/news/2016-05-link-tea-coffee-intake-barrett.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*