


# Poll: Many parents keep prescription opioids at home

16 May 2016, by Beata Mostafavi

**Figure 1. Proportion of parents who kept child's leftover pain medication at home**



Source: C.S. Mott Children's Hospital National Poll on Children's Health, 2016

Twice as many parents with no guidelines from their provider kept leftover pain pills at home compared to parents whose providers discussed what to do with them. Credit: C.S. Mott Children's Hospital National Poll on Children's Health

Nearly half of parents whose child had leftover pain medication from a surgery or illness say they kept the prescription opioids at home—representing a potential problem down the line.

Parents whose child's provider discussed what to do with the pills, however, were far more likely to dispose them properly, according to a report from the C.S. Mott Children's Hospital National Poll on Children's Health.


"We found that the amount of [pain medication](#) prescribed for children is frequently greater than the amount used, and too few parents recall clear direction from their provider about what to do with leftover medication," says Matthew M. Davis, M.D., director of the poll and professor of pediatrics and internal medicine at the University of Michigan's C.S. Mott Children's Hospital.

"This is a missed opportunity to prevent

prescription drug misuse among children. Many parents simply keep extra pain pills in their home. Those leftover pills represent easy access to narcotics for teens and their friends."

## Safe narcotic disposal

Overall, nearly a third of parents polled reported at least one pain medication prescription for their child, most often related to surgery, illness and injury. Most prescriptions (60 percent) were for narcotics, such as oxycodone or hydrocodone.



Half of parents with prescription painkillers for their child said they had leftover medication. Credit: University of Michigan Health System

Half of parents reported their child had leftover medication and only 8 percent returned it to the doctor or pharmacy. Another 47 percent kept the pills at home while 30 percent disposed of it in the trash or toilet, 6 percent used it for other family

members and 9 percent didn't remember what they did with them.

Twice as many parents (56 percent) with no guidelines from their provider kept leftover pain pills at home compared to parents whose providers discussed what to do with them. Only a third of parents said their child's provider gave instructions on disposing leftover pain medication.

"The epidemic of [prescription drug abuse](#) affects all sectors of the U.S. population, including children and teens," Davis says.

"For adolescents, a known point of access to narcotic pain medication is leftover pills from a prior prescription. Our poll suggests we need to do a much better job of explaining this risk to parents."

The Mott report comes just as the House of Representatives considers a series of new bills to tackle the opioid crisis that has led to major policy initiatives to limit access to narcotic pain medications. The Mott poll report presents findings from a nationally representative household survey conducted among 1,176 parents with at least one child aged 5 to 17.

#### **Takeaways for health care providers:**

- Providers cannot assume that parents understand the risks of easily accessible pain medication in the home. Providers need to explain this risk to parents, and offer guidance on appropriate options (e.g., return to provider office or pharmacy; safe disposal in trash, not flushing down the toilet).
- The poll results strongly suggest that when providers do offer direction to parents, they make better decisions with leftover medication.
- In this poll, half of parents whose children had received a prescription for pain medication had pills left over. This raises a fundamental question about how providers approach the task of prescribing pain medication to children: Are they prescribing "just enough" medication for a standard recovery, or additional doses "just in case"

there is prolonged pain?

- Providers should make careful decisions regarding the amount of narcotic pain medication prescribed to children, and parents should feel comfortable asking questions about the amount of medication prescribed.
- Parents need clearer guidance on all aspects of administering prescription medications for their children. This is particularly true for pain medication, which differs from the very specific manner in which other medications are handled; rather, the administration of pain medication is linked to the child's symptoms without a strict schedule or endpoint. Ideally, all [parents](#) should have guidance in this area

Provided by University of Michigan Health System

APA citation: Poll: Many parents keep prescription opioids at home (2016, May 16) retrieved 17 June 2022 from <https://medicalxpress.com/news/2016-05-poll-parents-prescription-opioids-home.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*