

Prophylactic antibiotics cut risk of anti-EGFR skin rash

1 June 2016



control arms (from 50 to 24 percent).

"Taking preemptive tetracyclines for several weeks at the start of anti-EGFR treatment can significantly reduce the incidence and severity of cutaneous acneiform rash," the authors write.

More information: [Abstract](#)
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

(HealthDay)—For patients receiving anti-epidermal growth factor receptor (EGFR) treatment, the risk of developing skin rash is reduced for those taking prophylactic antibiotics, according to a review published online May 23 in the *British Journal of Dermatology*.

Fausto Petrelli, M.D., from Piazzale Ospedale in Treviso, Italy, and colleagues conducted a systematic review and meta-analysis to examine whether [prophylactic antibiotics](#) can reduce the occurrence and severity of anti-EGFR drug-related skin rashes. Data were included for 1,073 [patients](#) from 13 studies.

The researchers found that, compared to patients without [antibiotic prophylaxis](#), patients in the prophylactic antibiotic arms had a reduced risk of developing a skin rash in all studies (odds ratio, 0.53). Based on 12 studies, moderate-to-severe toxicities were reduced by almost 70 percent (odds ratio, 0.32). This corresponded to a 26 percent absolute difference of high-grade skin rash versus

APA citation: Prophylactic antibiotics cut risk of anti-EGFR skin rash (2016, June 1) retrieved 27 October 2022 from <https://medicalxpress.com/news/2016-06-prophylactic-antibiotics-anti-egfr-skin-rash.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.