

Hypocaloric diet, exercise both reduce visceral adiposity

3 June 2016



More information: [Abstract](#)

[Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

(HealthDay)—Both hypocaloric diet and exercise reduce visceral adiposity (VAT), with exercise tending to have a larger effect on reducing VAT, according to a review published online May 23 in *Obesity Reviews*.

Rebecca J.H.M. Verheggen, M.D., from the Radboud University Medical Center in the Netherlands, and colleagues conducted a meta-analysis to examine the effects of [exercise](#) or [diet](#) on VAT. Data were included from 117 studies with 4,815 participants.

The researchers found that both exercise and diet caused VAT loss (P

"In conclusion, both exercise and diet reduce VAT. Despite a larger effect of diet on total body weight loss, exercise tends to have superior effects in reducing VAT," the authors write. "Finally, total body [weight loss](#) does not necessarily reflect changes in VAT and may represent a poor marker when evaluating benefits of lifestyle-interventions."

APA citation: Hypocaloric diet, exercise both reduce visceral adiposity (2016, June 3) retrieved 27 May 2022 from <https://medicalxpress.com/news/2016-06-hypocaloric-diet-visceral-adiposity.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.