

Vaginal pessary beneficial in pelvic organ prolapse

27 June 2016



"We provided further evidence in non-surgical treatment for POP," the authors write. "Prolapse symptoms and quality of life were improved in women using a vaginal pessary in addition to pelvic floor exercises."

More information: Full Text (subscription or payment may be required)

Copyright © 2016 HealthDay. All rights reserved.

(HealthDay)—For women with symptomatic pelvic organ prolapse (POP), use of vaginal pessary in addition to pelvic floor exercises is associated with improvements in prolapse symptoms and quality of life, according to a study published in the July issue of *Obstetrics & Gynecology*.

Rachel Y.K. Cheung, M.B.Ch.B., from the Chinese University of Hong Kong, and colleagues conducted a randomized trial involving women with symptomatic stage I to III POP. Participants were randomized to pelvic floor exercise training (control group; 128 women) or to pelvic floor exercise training and insertion of a vaginal pessary (pessary group; 132 women).

The researchers found that after 12 months, both groups had decreases in the Pelvic Organ Prolapse Distress Inventory of Pelvic Floor Distress Inventory and Pelvic Organ Prolapse Impact Questionnaire of Pelvic Floor Impact Questionnaire scores, with higher mean score differences in the pessary group (both P



APA citation: Vaginal pessary beneficial in pelvic organ prolapse (2016, June 27) retrieved 3 September 2022 from https://medicalxpress.com/news/2016-06-vaginal-pessary-beneficial-pelvic-prolapse.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.