

Smoking greatly reduces life expectancy for those with serious mental illness

11 August 2016, by Terri Mellow

Smokers with serious mental illness have their lives cut short by about 15 years, compared with people who have never smoked and who do not have serious mental illness, research from the University of Michigan shows.

They also die 10 years earlier than those with serious [mental illness](#) who have never smoked.

This means smoking may account for nearly two thirds of the overall difference in life expectancy between individuals with serious mental illness who smoke, and never smokers in the general population, says Jamie Tam, a doctoral candidate in the Department of Health Management and Policy at the U-M School of Public Health, who conducted the study.

"Smoking reduces life expectancy for everyone but we tend to underestimate the importance of smoking for people with comorbidities," Tam said. "We know from existing research that people with mental illness live shorter lives, but what wasn't known is how much of that is due to their mental illness and how much is due to the fact that many of them are smokers."

About 42 percent of people with serious [psychological distress](#) are smokers, compared with the [general population](#) at less than 20 percent.

To find out the contribution of smoking to reduced life expectancy among people with serious psychological distress, Tam and School of Public Health faculty members Kenneth Warner and Rafael Meza analyzed data from 328,000 individuals 25 years and older from the National Health Interview Survey gathered from 1997-2009. They linked it with the 2011 National Death Index.

Tam said a disproportionate number of those with serious mental illnesses—including bipolar disorder, schizophrenia and major depression—live in poverty, as they are often unemployed and many

face homelessness and social conditions that increase their likelihood of smoking and in turn their risk of other [health](#) conditions.

Deaths by suicide and accidents also occur more frequently in this population.

Among those with serious psychological distress, the researchers found that 47 percent of the deaths are due to the three leading health complications associated with smoking: heart disease, cancer and stroke.

"This highlights the need to prioritize preventive care, including smoking cessation programs, for people with mental illness," said Meza, senior author of the study and assistant professor of epidemiology at the U-M School of Public Health.

Many people with mental illness lack adequate health care—only 32 percent of those with serious psychological distress report seeing a mental health professional—and many don't get their physical health care needs met, Tam said.

"A large majority of people with mental illness don't access [mental health care](#), and even when they do, other health concerns like smoking are not addressed," she said. "There have been calls to integrate smoking interventions into mental [health care](#) but these recommendations haven't been fully implemented."

Their study appears in the *American Journal of Preventive Medicine*.

More information: Jamie Tam et al. Smoking and the Reduced Life Expectancy of Individuals With Serious Mental Illness, *American Journal of Preventive Medicine* (2016). DOI: [10.1016/j.amepre.2016.06.007](https://doi.org/10.1016/j.amepre.2016.06.007)

Provided by University of Michigan

APA citation: Smoking greatly reduces life expectancy for those with serious mental illness (2016, August 11) retrieved 19 August 2022 from <https://medicalxpress.com/news/2016-08-greatly-life-mental-illness.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.