

## Diabetes drug may also offer vascular protection

18 August 2016



Vincent DeMarco, Ph.D., is a research associate professor of endocrinology at the MU School of Medicine and lead author of the study. Credit: Justin Kelley, University of Missouri Health

Obesity and Type 2 diabetes are associated with vascular stiffening and the development of cardiovascular disease. Obese and diabetic premenopausal women are most at risk - even more than men of the same age who have similar health issues. A study by University of Missouri School of Medicine researchers found that a diabetes medication offered protection against arterial stiffness in overweight female mice, a finding that may have future implications for disease prevention in humans.

"The widespread overconsumption of a <u>western</u> <u>diet</u> high in fats and refined sugars is a contributing factor to the epidemic of obesity and diabetes around the world," said Vincent DeMarco, Ph.D., a research associate professor of endocrinology at the MU School of Medicine and lead author of the study. "Our previous studies showed that young female mice consuming a mostly western diet not only gained weight, but also exhibited arterial stiffening consistent with obese <u>premenopausal</u>

women. Our current study sought to understand what effects, if any, the diabetes medication linagliptin had on preventing vascular stiffness."

Linagliptin is a medication prescribed to lower blood glucose in patients with Type 2 diabetes. The medication works by blocking the enzyme dipeptidyl peptidase-4, or DPP-4. Previous studies have shown that the DPP-4 inhibitor also seemed to offer protection against vascular inflammation and oxidative stress - conditions associated with arterial stiffening.

DeMarco's team observed 34 <u>female mice</u> that were fed either a normal diet or a simulated western diet for four months. Another group of mice were fed a western diet containing a low dose of linagliptin. The team used an ultrasound system designed specifically for mice to evaluate stiffness of the aorta - the main artery that supplies blood to the circulatory system.

"The mice fed a western diet without receiving linagliptin gained weight and developed aortic stiffness," DeMarco said. "However, a big surprise to us was an almost total prevention of aortic stiffening in the mice that were fed the western diet along with linagliptin, even though this group gained as much weight as the other mice."

DeMarco cautions that more research is needed to determine if linagliptin could be used as a therapeutic tool in the future to prevent aortic stiffening and the cardiovascular risks associated with obesity and <u>diabetes</u>. Additionally, Linagliptin, like other DPP-4 inhibitors, can be expensive without insurance coverage.

"Based on the results of our study, it is tempting to speculate that linagliptin could target <u>arterial</u> <u>stiffness</u> and reduce the risk of <u>cardiovascular</u> <u>disease</u>," DeMarco said. "However, results from clinical trials already in progress will be needed to determine what, if any, future role linagliptin could



play in the management of obesity-related cardiovascular disease."

**More information:** Camila Manrique et al, Dipeptidyl peptidase-4 inhibition with linagliptin prevents western diet-induced vascular abnormalities in female mice, *Cardiovascular Diabetology* (2016). <u>DOI:</u> <u>10.1186/s12933-016-0414-5</u>

Provided by University of Missouri-Columbia APA citation: Diabetes drug may also offer vascular protection (2016, August 18) retrieved 24 September 2022 from <u>https://medicalxpress.com/news/2016-08-diabetes-drug-vascular.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.