

Albuminuria linked to higher nighttime SBP in hypertension

22 August 2016



nighttime SBP, particularly in those with diabetes with very high albuminuria and low eGFR," the authors write.

The Spanish Ambulatory BP Monitoring Registry cohort is maintained by an unrestricted grant from Lacer Laboratorie.

More information: [Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

(HealthDay)—For patients with hypertension, albuminuria is associated with higher nighttime systolic blood pressure (SBP), according to a study published online Aug. 11 in *Diabetes Care*.

Gema Ruiz-Hurtado, Ph.D., from the Hospital Universitario 12 de Octubre in Madrid, and colleagues examined the quantitative differences in nighttime SBP across albuminuria levels in patients with and without [diabetes](#) and [chronic kidney disease](#). Data were analyzed for 16,546 patients from the Spanish Ambulatory BP Monitoring Registry cohort. Patients were classified according to estimated [glomerular filtration rate](#) (eGFR) and urine albumin-to-creatinine ratio.

The researchers observed a statistically significant and clinically substantial higher nighttime SBP for high albuminuria (6.8 mm Hg higher than with normoalbuminuria; P

"Albuminuria in hypertensive [patients](#) is accompanied by quantitatively striking higher

APA citation: Albuminuria linked to higher nighttime SBP in hypertension (2016, August 22) retrieved 29 April 2021 from <https://medicalxpress.com/news/2016-08-albuminuria-linked-higher-nighttime-sbp.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.