

Playing with sports concussion doubles recovery time: study

August 29 2016, by Lindsey Tanner

Continuing to play despite a concussion doubles recovery time for teen athletes and leads to worse short-term mental function than in those immediately removed from action, a study found.

It's billed as the first to compare recovery outcomes for athletes removed from a game or practice compared with those who aren't. The study was small, involving 69 teens treated at a University of Pittsburgh Medical Center concussion clinic, but the results bolster evidence supporting the growing number of return-to-play laws and policies nationwide

The study was published Monday in the journal *Pediatrics*.

KEEPING SCORE

The study involved athletes aged 15 on average from several sports, including football, soccer, ice hockey and basketball who had concussions during a game or practice. Half continued to play and took 44 days on average to recover from symptoms, versus 22 days in those who were immediately sidelined.

Sidelined players reported symptoms immediately, including dizziness, headaches, mental foginess and fatigue, and were diagnosed with concussions by trainers or team physicians. The others, who continued playing for 19 minutes on average, delayed reporting symptoms and were diagnosed later.

Those who continued to play had worse scores on [mental function](#) tests performed eight days after the concussion and 30 days after the concussion. Medical records showed mental function had been similar in all players before their concussions.

RISKY RETURNS

Return-to-play policies are widespread, especially in youth athletics, and they typically recommend sidelining players after a suspected concussion until symptoms resolve. One of the main reasons is to prevent a rare condition called second-impact syndrome—potentially fatal brain swelling or bleeding that can occur when a player still recovering from a concussion gets hit again in the head.

The study results show that a prolonged recovery is another important risk from returning to play too soon—one that "no one had really calculated" until now, said Dr. Allen Sills, a Vanderbilt University neurosurgeon. He was not involved in the research.

NOT REPORTED

About 300,000 sports-related concussions occur each year nationwide among all ages. In high school athletics, they occur at a rate of almost 3 per 10,000 games or practices.

Evidence suggests up to 50 percent of concussions in teen sports aren't reported. Athletes are sometimes not aware they've experienced a concussion, or they suspect a head injury but continue playing because "they don't want to let their teammates down," said University of Arkansas [concussion](#) researcher R.J. Elbin, the study's lead author.

The results "give us more ammunition" to persuade [young athletes](#) to heed the return-to-play advice, Elbin said.

More information: *Pediatrics*: www.pediatrics.org

CDC: www.cdc.gov/headsup/youthsports/

© 2016 The Associated Press. All rights reserved.

Citation: Playing with sports concussion doubles recovery time: study (2016, August 29)
retrieved 17 December 2022 from
<https://medicalxpress.com/news/2016-08-sports-concussion-recovery.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.