

Why bones don't heal: Researchers identify risk factors for nonunion of fractures

September 7 2016

Dr. Robert Zura, the Robert D'Ambrosia Professor and Head of Orthopaedic Surgery at LSU Health New Orleans School of Medicine, was part of a research team that identified risk factors which may help orthopaedic surgeons better predict a serious complication of bone fractures. Fracture nonunion may be increasing as more patients survive serious fractures. The paper was published September 7, 2016, in the Online First section of *JAMA Surgery*.

To better understand the risk factors that contribute to the failure of [fractured bones](#) to heal, a national team of orthopaedic surgeons examined health records from 90.1 million patients. Their goal was to describe the epidemiology of fracture nonunion in adults using the information available to physicians at patient presentation. They focused on patients 18-63 years old, representing 6,725 patients with 309,330 fractures. They researched risk factors likely to be of general concern to [orthopaedic surgeons](#) and pooled data to identify 45 variables of interest.

The researchers found a nonunion rate of 4.93% overall with substantial variation from bone to bone. The lowest nonunion bone rates were in metacarpal at 1.47% and radius at 2.10%. The highest nonunion rates were in scaphoid at 15.46%, tibia+fibula at 13.95%, and femur at 13.86%.

The presence of other diseases boosted risk. Osteoarthritis, rheumatoid arthritis, and diabetes increased the odds of nonunion by at least 40%.

Use of certain medications such as analgesics also upped risk. Other medications were protective. Antidiabetic medications, other than insulin, and oral contraceptives are inversely associated with nonunion.

Nonunion odds were significantly increased by many risk factors including the number of fractures, use of prescription analgesics (NSAIDs+opioids), operative treatment, open fracture, anticoagulant use, osteoarthritis with or without rheumatoid arthritis, anticonvulsant use with or without benzodiazepine, opioid use, diabetes, high-energy injury, osteoporosis, male gender, insulin use, smoking, obesity, antibiotic use, Vitamin D deficiency, diuretic use and kidney insufficiency.

The researchers contributed novel insights into the causes of fracture nonunion. They identified the most significant risk factors - the severity and location of the fracture, the presence of other diseases and medication use. This information could be used to develop a predictive algorithm.

"Understanding key risk factors and their interplay will help us determine which fractures are at greatest risk of nonunion," concludes Dr. Robert Zura, Professor and Head of Orthopaedics at LSU Health New Orleans. "We can alter our approach to those and modify those [risk factors](#) we can, such as prescription medicines, to improve healing."

More information: *JAMA Surgery*,
<http://archsurg.jamanetwork.com/article.aspx?articleid=2547685>

Provided by Louisiana State University

Citation: Why bones don't heal: Researchers identify risk factors for nonunion of fractures

(2016, September 7) retrieved 2 July 2023 from <https://medicalxpress.com/news/2016-09-bones-dont-factors-nonunion-fractures.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.