


High variability suggests glycemic index is unreliable indicator of blood sugar response

September 8 2016



Each blue dot represents the average of three glycemic index value determinations for an individual subject. Horizontal bars represent the standard deviations. Adapted from Matthan et al (2016). Credit: Tufts University

The glycemic index of a given food, a value that aims to quantify how fast blood sugar rises after eating it, can vary by an average of 20 percent within an individual and 25 percent among individuals, report scientists from the Jean Mayer USDA Human Nutrition Research Center on Aging (USDA HRNCA) at Tufts University.

In randomized, controlled, repeated tests involving 63 healthy adults, researchers found that individual [blood sugar](#) responses after consuming a fixed amount of white bread could range across all three [glycemic index](#) categories (low, medium, or high). Part of this variability could be attributed to insulin index and baseline HbA1c levels, which reflect long-term glucose control—evidence that glycemic index values are influenced by an individual's metabolic responses to food.

The study, published in the *American Journal of Clinical Nutrition* on Sept. 7, suggests glycemic index has limited utility as a tool to predict how a food affects blood sugar levels.

"Glycemic index values appear to be an unreliable indicator even under highly standardized conditions, and are unlikely to be useful in guiding food choices," said lead study author Nirupa Matthan, Ph.D., scientist in the Cardiovascular Nutrition Laboratory at the USDA HNRCA. "If someone eats the same amount of the same food three times, their blood glucose response should be similar each time, but that was not observed in our study. A food that is [low glycemic index](#) for you one time you eat it could be high the next time, and it may have no impact on blood sugar for me."

Developed as a way to help diabetic individuals control their blood sugar, glycemic index is intended to represent the inherent effect a food has on blood sugar levels. However, glycemic index is becoming used for broader purposes such as food labeling, and has served as the basis for several popular diets.

To study whether glycemic index values are accurate and reproducible, Matthan and her colleagues recruited 63 volunteers, who underwent six testing sessions over 12 weeks. Volunteers fasted and abstained from exercise and alcohol before each session. They then consumed either white bread, a simple carbohydrate that served as the test food, or a glucose drink, which served as a reference control, in random order. Each contained 50 grams of available carbohydrate. Blood glucose levels were measured at multiple time points for five hours after eating, and glycemic index was calculated by standard formulas.

An unreliable guide

The team found that the average glycemic index value of white bread for the study population was 62, placing it in the category of a "medium" glycemic index food.

However, deviations averaged 15 points in either direction, effectively placing white bread in all three glycemic index categories. It would be considered a low glycemic index food (average values of 35 to 55) for 22 of the volunteers, intermediate glycemic index (57 to 67) for 23 volunteers, and [high glycemic index](#) (70 to 103) for 18 volunteers. Even within the same individual, glycemic index values could differ by more than 60 points between trials.

"Reports frequently tout the benefits of choosing foods with low glycemic index and glycemic load values. Our data suggest those values may not be reliable in terms of a daily intake. A better approach to choosing foods is to consume a diet primarily composed of vegetables, fruits, whole grains, nonfat and low-fat dairy products, fish, legumes (beans), lean meats with preference to preparing food with liquid vegetable oils, and equally as important, to choose healthy foods and beverages you really enjoy," said senior study author Alice H. Lichtenstein, D.Sc., senior scientist and director of the Cardiovascular

Nutrition Laboratory at the USDA HNRCA. Lichtenstein is also the Gershoff Professor at the Friedman School of Nutrition Science and Policy at Tufts.

The variability in glycemic index values occurred despite sample sizes larger than required by standard calculations. The study cohort of 63 individuals far exceeded the 10 individuals used by typical glycemic index methodology, as did the six feeding challenges and five-hour blood glucose measuring window.

The team also tested for the influence of biological characteristics: sex, body-mass index, blood pressure, physical activity, and several others. Most factors had only a minor statistical effect on glycemic index variability. Blood insulin response as measured by insulin index and HbA1c, a measure of longer term glucose control, had the largest effect, accounting for 15 and 16 percent of the variability, respectively.

The authors note their findings do not suggest that a high glycemic index food may be healthy, or that a low glycemic index food unhealthy. Both glycemic index and glycemic load—a value that adjusts glycemic index to serving size—reflect only food containing carbohydrates, and no one eats food in isolation. The high variability of glycemic index and glycemic load reveal limitations in their clinical and public health applicability, and glycemic index estimates and subsequent associations with chronic disease risk needs to be reconsidered, say the authors.

"Based on our results, we feel strongly that glycemic index is impractical for use in [food](#) labeling or in dietary guidelines at the individual level," Matthan said. "If your doctor told you your LDL cholesterol value could vary by 20 percent, it would be the difference between being normal or at high risk for heart disease. I don't think many people would find that acceptable."

More information: N. R. Matthan et al. Estimating the reliability of glycemic index values and potential sources of methodological and biological variability, *American Journal of Clinical Nutrition* (2016). DOI: [10.3945/ajcn.116.137208](https://doi.org/10.3945/ajcn.116.137208)

Provided by Tufts University

Citation: High variability suggests glycemic index is unreliable indicator of blood sugar response (2016, September 8) retrieved 14 May 2023 from <https://medicalxpress.com/news/2016-09-high-variability-glycemic-index-unreliable.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.