

# High glycemic index diet raises hepatic fat, glycogen stores

9 September 2016



"Compared to an LGI diet, a one-week HGI diet increased hepatic fat and glycogen stores," the authors write. "This may have important clinical relevance for dietary interventions in the prevention and management of non-alcoholic fatty liver disease."

Several authors disclosed financial ties to Unilever.

**More information:** [Abstract](#)  
[Full Text \(subscription or payment may be required\)](#)

Copyright © 2016 [HealthDay](#). All rights reserved.

(HealthDay)—A one-week high glycemic index (HGI) diet increases hepatic fat and glycogen stores in healthy adults compared with a low glycemic index (LGI) diet, according to a study published online Sept. 4 in *Diabetes, Obesity and Metabolism*.

Stephen Bawden, Ph.D., from the University of Nottingham in the United Kingdom, and colleagues examined the acute and long-term effects of LGI and HGI diets in healthy volunteers. Eight healthy males underwent testing before and after a seven-day macronutrient and energy matched HGI or LGI [diet](#), followed by a minimum four-week washout period, and repetition of the intervention with the alternative diet.

The researchers found that following the HGI versus LGI test meal, there were significantly greater plasma glucose and insulin peak values and area under the curve. Following the HGI test meal, there was a greater increase in hepatic glycogen concentrations (P intervention).

APA citation: High glycemic index diet raises hepatic fat, glycogen stores (2016, September 9) retrieved 4 May 2021 from <https://medicalxpress.com/news/2016-09-high-glycemic-index-diet-hepatic.html>

*This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.*