

Occipital nerve stimulation effective for chronic migraine

29 October 2016



of the occipital nerves in this single-center trial," the authors write.

More information: Full Text (subscription or payment may be required)

Copyright © 2016 HealthDay. All rights reserved.

(HealthDay)—For patients with chronic migraine (CM), peripheral nerve stimulation of the occipital nerves reduces the number of headache days, according to a study published online Oct. 25 in *Pain Practice*.

Nagy A. Mekhail, M.D., Ph.D., from the Cleveland Clinic, and colleagues implanted 20 <u>patients</u> at a single center with a neurostimulation system, and randomized them to an active or control group for 12 weeks. Patients received open-label treatment for an additional 40 weeks.

The researchers observed a reduction in the number of headache days per month (8.51 days; P pain intensity. All patients had reductions in Migraine Disability Assessment and Zung Pain and Distress scores. At least one adverse event was reported by 15 of the patients, with a total of 20 adverse events reported.

"Our results support the 12-month efficacy of 20 CM patients receiving peripheral nerve stimulation



APA citation: Occipital nerve stimulation effective for chronic migraine (2016, October 29) retrieved 8 May 2021 from https://medicalxpress.com/news/2016-10-occipital-nerve-effective-chronic-migraine.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.