

Tobacco linked to 40 percent of US cancers

November 10 2016



Credit: Vera Kratochvil/public domain

Tobacco use remains the most preventable cause of cancer, and 40 percent of diagnosed US cancer cases may have a link to its use, health authorities said Thursday.

Lung cancer, [acute myeloid leukemia](#), and tumors of the mouth and throat, [voice box](#), esophagus, stomach, kidney, pancreas, liver, bladder, cervix, colon and rectum are all caused by tobacco use, according the report by the US Centers for Disease Control and Prevention.

"There are more than 36 million smokers in the US," said CDC director Tom Frieden in a statement.

"Sadly, nearly half could die prematurely from tobacco-related illnesses, including six million from cancer, unless we implement the programs that will help smokers quit."

The CDC Vital Signs report found that every year from 2009 to 2013, about 660,000 people in the United States were diagnosed with a cancer related to tobacco use.

About 343,000 people died each year from these cancers.

"Three in ten cancer deaths were due to cigarette smoking," said the report.

Lung cancer is the top cause of tobacco related smoking, followed by colon cancer and pancreatic cancer.

Cigarette-smoking is on the decline, down from 21 percent (45.1 million) of the US population in 2005 to 15 percent (36.5 million) in 2015.

This is the lowest point for cigarette smoking since data began to be collected in 1965.

Since 1990, about 1.3 million tobacco-related cancer deaths have been avoided, the CDC added.

© 2016 AFP

Citation: Tobacco linked to 40 percent of US cancers (2016, November 10) retrieved 30 March 2023 from <https://medicalxpress.com/news/2016-11-tobacco-linked-percent-cancers.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.